

FCRM PTO-1390

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE  
TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

ATTORNEY DOCKET NUMBER  
JANS-0004

U.S. APPLICATION NO. (if known see 37 C.F.R. 1.5)

09/868535

INTERNATIONAL APPLICATION NO.  
PCT/EP99/10176

INTERNATIONAL FILING DATE  
15 December 1999

PRIORITY DATE CLAIMED  
19 December 1998

TITLE OF INVENTION ANTIHISTAMINIC SPIRO COMPOUNDS

APPLICANT(S) FOR DO/EO/US Frans Eduard JANSSEN and Joseph Elisabeth LEENAERTS

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
  - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☒ has been transmitted by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
  - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) 35 U.S.C. 371(c)(4).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:


11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.  
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:
  - A copy of the Published PCT Application by WIPO under No. WO 00/37470, including the Search Report and references cited.
  - A copy of the International Preliminary Examination Report; Associate Power of Attorney.

EXPRESS MAIL Mailing Label No. EL 922205782 US  
Date of Deposit: June 19, 2001

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231

MAILER John Hill

SIGNATURE John Hill

U.S. APPLICATION NO. (if known) <b>09/868535</b> (R. 1.5)		INTERNATIONAL APPLICATION NO. <b>PCT/EP99/10176</b>		ATTORNEY DOCKET NUMBER <b>JANS-0004</b>	
17. <u>  </u> The following fees are submitted: <b>Basic National Fee (37 CFR 1.492(a)(1) - (5)):</b> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO..... <b>\$1,000.00</b>  International preliminary examination fee (37 CFR 1.482 not paid to USPTO but International Search Report has been prepared by the EPO or JPO..... <b>\$860.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO..... <b>\$710.00</b>  International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4)..... <b>\$690.00</b>  International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4)..... <b>\$100.00</b>  <div style="text-align: right;"><b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b></div>				<u>CALCULATIONS</u> <u>PTO USE ONLY</u>	
Surcharge of \$130.00 for furnishing the oath or declaration later than <u>  </u> 20 <u>  </u> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
Claims	Number Filed	Number Extra	Rate		
Total claims	15 - 20 =	0	X \$18.00	\$	
Independent Claims	1 - 3 =	0	x \$80.00	\$	
Multiple dependent claims(s) (if applicable)			+ \$270.00	\$	
<b>TOTAL OF ABOVE CALCULATIONS =</b>				<b>\$860.00</b>	
Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by ½.				\$	
<b>SUBTOTAL =</b>				<b>\$860.00</b>	
Processing fee of <b>\$130.00</b> for furnishing the English translation later than <u>  </u> 20 <u>  </u> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				+	
<b>TOTAL NATIONAL FEE =</b>				<b>\$860.00</b>	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). <b>\$40.00</b> per property				+	
<b>TOTAL FEES ENCLOSED =</b>				<b>\$860.00</b>	
				<b>Amount to be:</b>	
				<b>refunded</b>	\$
				<b>charged</b>	\$
a. <input checked="" type="checkbox"/> A check in the amount of <b>\$ 860.00</b> to cover the above fee is enclosed.  b. <u>  </u> Please charge my Deposit Account No. 23-3050 in the amount of \$ <u>      </u> to cover the above fees. A duplicate copy of this sheet is enclosed.  c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 23-3050. A duplicate copy of this sheet is enclosed.					
<b>NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must          be filed and granted to restore the application to pending status.</b>					
SEND ALL CORRESPONDENCE TO: David N. Farsiou Woodcock Washburn Kurtz Mackiewicz & Norris LLP One Liberty Place - 46th Floor Philadelphia, PA 19103 (215) 568-3100			 SIGNATURE  <b>David N. Farsiou</b> NAME  <b>44,104</b> REGISTRATION NUMBER		

DOCKET NO.: JANS-0004/JAB-1442

PATENT

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re patent application of:

Janssens et al.

U.S. Serial No.: Not yet assigned

Group Art Unit: Not yet assigned

Filed: Not yet assigned

Examiner: Not yet assigned

For: ANTIHISTAMINIC SPIRO COMPOUNDS

Assistant Commissioner of  
Patents & Trademarks  
Washington, DC 20231

PRELIMINARY AMENDMENT


Please amend the application, without prejudice, as follows.

In the Claims

Please cancel claim 9.

Please amend claims 1, 4-8, 10-12, and 14-15 as follows:

1. (Amended) A compound of formula



or a prodrug, a *N*-oxide, an addition salt, a quaternary amine or a stereochemically isomeric form thereof wherein

$R^1$  is hydrogen,  $C_{1-6}$ alkyl, halo, formyl, carboxyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylcarbonyl,  $N(R^3R^4)C(=O)-$ ,  $N(R^3R^4)C(=O)N(R^5)-$ , ethenyl substituted with carboxyl or  $C_{1-6}$ alkyloxycarbonyl, or  $C_{1-6}$ alkyl substituted with hydroxy, carboxyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $N(R^3R^4)C(=O)-$ ,  $C_{1-6}$ alkyl $C(=O)N(R^5)-$ ,  $C_{1-6}$ alkyl $S(=O)_2N(R^5)-$  or  $N(R^3R^4)C(=O)N(R^5)-$ ;

wherein each  $R^3$  and each  $R^4$  independently are hydrogen or  $C_{1-4}$ alkyl;

$R^5$  is hydrogen or hydroxy;

$R^2$  is hydrogen,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy $C_{1-6}$ alkyl,  $N(R^3R^4)C(=O)-$ , aryl or halo;

$n$  is 1 or 2;

-A-B- represents a bivalent radical of formula

-Y-CH=CH- (a-1);

-CH=CH-Y- (a-2); or

-CH=CH-CH=CH- (a-3);

wherein each hydrogen atom in the radicals (a-1) to (a-3) may independently be replaced by  $R^6$  wherein  $R^6$  is selected from  $C_{1-6}$ alkyl, halo, hydroxy,  $C_{1-6}$ alkyloxy, ethenyl substituted with carboxyl or  $C_{1-6}$ alkyloxycarbonyl, hydroxy $C_{1-6}$ alkyl, formyl, carboxyl or hydroxycarbonyl $C_{1-6}$ alkyl;

each Y independently is a bivalent radical of formula -O-, -S- or -NR<sup>7</sup>-;

wherein  $R^7$  is hydrogen,  $C_{1-6}$ alkyl or  $C_{1-6}$ alkylcarbonyl;

Z is a bivalent radical of formula

-(CH<sub>2</sub>)<sub>p</sub>- (b-1),

-CH=CH- (b-2),

-CH<sub>2</sub>-CHOH- (b-3),

-CH<sub>2</sub>-O- (b-4),

-CH<sub>2</sub>-C(=O)- (b-5), or


-CH<sub>2</sub>-C(=NOH)- (b-6),

with the proviso that the bivalent radicals (b-3), (b-4), (b-5) and (b-6) are connected to the nitrogen of the imidazole ring via their -CH<sub>2</sub>- moiety;

wherein  $p$  is 1, 2, 3 or 4;

L is hydrogen;  $C_{1-6}$ alkyl;  $C_{2-6}$ alkenyl;  $C_{1-6}$ alkylcarbonyl;  $C_{1-6}$ alkyloxycarbonyl;  $C_{1-6}$ alkyl substituted with one or more substituents each independently selected from hydroxy, carboxyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl, aryl, aryloxy, cyano or  $R^8HN-$  wherein  $R^8$  is hydrogen,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylcarbonyl; or

L represents a radical of formula




Alk represents  $\text{C}_{1-4}$  alkanediyl ;

Y represents O, S or NH ;

Het<sup>1</sup> and Het<sup>2</sup> each represent furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl or imidazolyl each optionally substituted with one or two  $\text{C}_{1-4}$  alkyl substituents; pyrrolyl or pyrazolyl optionally substituted with formyl, hydroxy $\text{C}_{1-4}$ alkyl, hydroxycarbonyl,  $\text{C}_{1-4}$ alkyloxy-carbonyl or with one or two  $\text{C}_{1-4}$  alkyl substituents; thiadiazolyl or oxadiazolyl optionally substituted with amino or  $\text{C}_{1-4}$ alkyl; pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl each optionally substituted with  $\text{C}_{1-4}$ alkyl,  $\text{C}_{1-4}$ alkyloxy, amino, hydroxy or halo; and

Het<sup>3</sup> represents furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl or imidazolyl each optionally substituted with one or two  $\text{C}_{1-4}$  alkyl substituents; pyrrolyl or pyrazolyl optionally substituted with formyl, hydroxy $\text{C}_{1-4}$ alkyl, hydroxycarbonyl,  $\text{C}_{1-4}$ alkyloxy-carbonyl or with one or two  $\text{C}_{1-4}$  alkyl substituents; thiadiazolyl or oxadiazolyl optionally substituted with amino or  $\text{C}_{1-4}$ alkyl; pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl each optionally substituted with  $\text{C}_{1-4}$ alkyl,  $\text{C}_{1-4}$ alkyloxy, amino, hydroxy, halo, 4,5-dihydro-5-oxo-1H-tetrazolyl substituted with  $\text{C}_{1-4}$ alkyl, 2-oxo-3-oxazolidinyl, 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl or a radical of formula



wherein

A-Z represents  $\text{S}-\text{CH}=\text{CH}$ ,  $\text{S}-\text{CH}_2-\text{CH}_2$ ,  $\text{S}-\text{CH}_2-\text{CH}_2-\text{CH}_2$ ,  $\text{CH}=\text{CH}-\text{CH}=\text{CH}$ , or  $\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2$ ;


aryl is phenyl or phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, hydroxy,  $\text{C}_{1-4}$ alkyl, polyhalo $\text{C}_{1-4}$ alkyl, cyano, aminocarbonyl,  $\text{C}_{1-4}$ alkyloxy or polyhalo $\text{C}_{1-4}$ alkyloxy;

with the proviso that 5,6-dihydrospiro[imidazo[1,2-b][3]benzazepine-11[11H],4'-piperidine] and pharmaceutically acceptable addition salts thereof are not included.

4. (Amended) A compound according to claim 1 wherein -A-B- is a bivalent radical of formula  $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$  (a-3) or  $-\text{CH}=\text{CH}-\text{Y}-$  (a-2).


5. (Amended) A compound according to claim 1 wherein Z is  $-(CH_2)_p-$  (b-1),  $-CH=CH-$  (b-2), or  $-CH_2-O-$  (b-4).
6. (Amended) A compound according to claim 1, wherein L is hydrogen,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl, carboxy $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxycarbonyl, or  $C_{1-6}$ alkyloxycarbonyl $C_{1-6}$ alkyl.
7. (Amended) A compound according to claim 1 wherein  $R^1$  is hydroxy $C_{1-6}$ alkyl, formyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkyloxy $C_{1-6}$ alkyl,  $N(R^3R^4)C(=O)-$ , halo or hydrogen.
8. (Amended) A compound according to claim 1 wherein the compound is
  - 5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-carboxamide dihydrochloride;
  - 1'-butyl-5,6-dihydrospiro[imidazo[2,1-*b*] [3] benzazepine-11-[11*H*],4'-piperidine];
  - 6,11-dihydro-1'-methylspiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine] cyclohexylsulfamate(1:2);
  - 6,11-dihydrospiro[5-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-methanol] (E)-2-butenedioate (2:1);
  - 3-chloro-6,11-dihydrospiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine] (E)-2-butenedioate (1:1);
  - 6,11-dihydro-3-(methoxymethyl)spiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]. (E)-2-butenedioate (1:1);
  - 6,11-dihydro-1'-(2-hydroxyethyl)spiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-carboxamide;
  - 6,11-dihydro-1'-methylspiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-carboxamide monohydrate;
  - ethyl 3-(aminocarbonyl)-6,11-dihydro- $\alpha$ -phenylspiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-propanoate monohydrochloride;
  - 3-(aminocarbonyl)-6,11-dihydrospiro[5*H*-imidazo[2,1-*b*][3]-benzazepine-11,4'-piperidine]-1'-carboxylate;
  - spiro[10*H*-imidazo[1,2-*a*]thieno[3,2-*d*]azepine-10,4'-piperidine];
  - 6,11-dihydrospiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-2,3-dicarboxamide dihydrochloride monohydrate; or
  - a prodrug, a *N*-oxide, an addition salt, a quaternary amine or a stereochemically isomeric form thereof.

10. (Amended) A pharmaceutical composition comprising a pharmaceutically acceptable carrier, and as active ingredient a therapeutically effective amount of a compound as defined in claim 1.
11. (Amended) A process of preparing a composition as claimed in claim 10, wherein a pharmaceutically acceptable carrier is mixed with a therapeutically effective amount of a compound as defined in claim 1.
12. (Amended) A compound of formula




or a *N*-oxide, an addition salt, a quaternary amine or a stereochemically isomeric form thereof wherein P is a protective group and n, -A-B-, Z, R<sup>1</sup> and R<sup>2</sup> are defined as in claim 1, with the proviso that 6,11-dihydro-1'-(phenylmethyl)-5*H*-spiro[imidazo[1,2-*b*][3]-benzazepine-11,4'-piperidine] (E)-2-butenedioate(1:2) is not included.

14. (Amended) A process of preparing a compound as claimed in claim 1, comprising
- deprotecting an intermediate of formula (II), followed optionally by derivatizing either the piperidine moiety, or the imidazole moiety, or both the piperidine moiety and the imidazole moiety




with P being a protective group;

- derivatizing an intermediate of formula (II) at the imidazole moiety, to form an intermediate of formula (II-a), followed by deprotecting the piperidine moiety, and followed optionally by derivatizing the piperidine moiety




c) cyclizing an intermediate of formula (III) in the presence of an appropriate acid, to form a compound of formula (I-a)



and, optionally, converting compounds of formula (I) and (I-a) into each other, and further, optionally, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid; and, optionally, preparing stereochemically isomeric forms or N-oxide forms thereof.


15. (Amended) A process of preparing a compound as claimed in claim 13, comprising,
  - a) cyclizing a compound of formula (XI) with an appropriate acid, to form a compound of formula (II-k), followed optionally by derivatizing the imidazole moiety, to form a compound of formula (II-a-1)





with  $Z_1$  being a bivalent radical of formula  $-(CH_2)_p-$ , wherein  $p$  is 1, 2, 3 or 4; and

- b) reacting a tricyclic moiety of formula (XIV) with a reagent of formula (XV) under an inert atmosphere in a reaction inert solvent in the presence of a suitable base, to form a compound of formula (II-l), followed optionally by derivatizing the imidazole moiety to form a compound of formula (II-a-2)



with  $W_5$  being a suitable leaving group, and  $Z_2$  being a bivalent radical of formula  $-(CH_2)_p-$ , or  $-CH_2-O-$ , wherein  $p$  is 1, 2, 3 or 4.

REMARKS

Claims 1-8 and 10-15 are presented. Claims 1, 4-8, 10-12, and 14-15 have been amended. Claim 9 has been canceled and no claims have been added.

The claims have been amended to remove multiple dependencies and to effect minor editorial amendments and are not intended to alter the scope or meaning of the original claims.

Attached hereto is a marked-up version of the changes made to the specification and claims by the current amendment. The attached page is captioned "Version with markings to show changes made." Entrance of the foregoing amendments and an early and favorable Action is respectfully requested.

Respectfully submitted,

Date: 6/19/01



David N. Farsiou  
Registration No. 44,104

WOODCOCK WASHBURN KURTZ  
MACKIEWICZ & NORRIS  
One Liberty Place - 46th Floor  
Philadelphia, PA 19103  
(215) 568-3100

DOCKET NO.: JANS-0004/JAB-1442


## VERSION WITH MARKINGS TO SHOW CHANGES MADE

## In the Claims:

Please cancel claim 9.

Please amend claims 1, 4-8, 10-12, and 14-15 as follows:

1. (Amended) A compound of formula



or a prodrug, a *N*-oxide, an addition salt, a quaternary amine or a stereochemically isomeric form thereof wherein

$R^1$  is hydrogen,  $C_{1-6}$ alkyl, halo, formyl, carboxyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylcarbonyl,  $N(R^3R^4)C(=O)-$ ,  $N(R^3R^4)C(=O)N(R^5)-$ , ethenyl substituted with carboxyl or  $C_{1-6}$ alkyloxycarbonyl, or  $C_{1-6}$ alkyl substituted with hydroxy, carboxyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $N(R^3R^4)C(=O)-$ ,  $C_{1-6}$ alkyl $C(=O)N(R^5)-$ ,  $C_{1-6}$ alkyl $S(=O)_2N(R^5)-$  or  $N(R^3R^4)C(=O)N(R^5)-$ ;

wherein each  $R^3$  and each  $R^4$  independently are hydrogen or  $C_{1-4}$ alkyl;

$R^5$  is hydrogen or hydroxy;

$R^2$  is hydrogen,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy $C_{1-6}$ alkyl,  $N(R^3R^4)C(=O)-$ , aryl or halo;

$n$  is 1 or 2;

-A-B- represents a bivalent radical of formula

-Y-CH=CH- (a-1);

-CH=CH-Y- (a-2); or

-CH=CH-CH=CH- (a-3);

wherein each hydrogen atom in the radicals (a-1) to (a-3) may independently be replaced by  $R^6$

wherein  $R^6$  is selected from  $C_{1-6}$ alkyl, halo, hydroxy,  $C_{1-6}$ alkyloxy, ethenyl substituted with carboxyl or  $C_{1-6}$ alkyloxycarbonyl, hydroxy $C_{1-6}$ alkyl, formyl, carboxyl [and] or hydroxycarbonyl $C_{1-6}$ alkyl;

each Y independently is a bivalent radical of formula -O-, -S- or -NR<sup>7</sup>-;

wherein  $R^7$  is hydrogen,  $C_{1-6}$ alkyl or  $C_{1-6}$ alkylcarbonyl;

Z is [is] a bivalent radical of formula

$-(CH_2)_p-$  (b-1),  
 $-CH=CH-$  (b-2),  
 $-CH_2-CHOH-$  (b-3),  
 $-CH_2-O-$  (b-4),  
 $-CH_2-C(=O)-$  (b-5), or  
 $-CH_2-C(=NOH)-$  (b-6),

[provided] with the proviso that the bivalent radicals (b-3), (b-4), (b-5) and (b-6) are connected to the nitrogen of the imidazole ring via their  $-CH_2-$  moiety;

wherein p is 1, 2, 3 or 4;

L is hydrogen;  $C_{1-6}$ alkyl;  $C_{2-6}$ alkenyl;  $C_{1-6}$ alkylcarbonyl;  $C_{1-6}$ alkyloxycarbonyl;  $C_{1-6}$ alkyl substituted with one or more substituents each independently selected from hydroxy, carboxyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl, aryl, aryloxy, cyano or  $R^8HN-$  wherein  $R^8$  is hydrogen,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylcarbonyl; or

L represents a radical of formula

$-Alk-Y-Het^1$  (c-1),  
 $-Alk-NH-CO-Het^2$  (c-2) or  
 $-Alk-Het^3$  (c-3) ; wherein


Alk represents  $C_{1-4}$ alkanediyl ;

Y represents O, S or NH ;

Het<sup>1</sup> [,] and Het<sup>2</sup> [and Het<sup>3</sup>] each represent furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl or imidazolyl each optionally substituted with one or two  $C_{1-4}$ alkyl substituents; pyrrolyl or pyrazolyl optionally substituted with formyl, hydroxy $C_{1-4}$ alkyl, hydroxycarbonyl,  $C_{1-4}$ alkyloxycarbonyl or with one or two  $C_{1-4}$ alkyl substituents; thiadiazolyl or oxadiazolyl optionally substituted with amino or  $C_{1-4}$ alkyl; pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl each optionally substituted with  $C_{1-4}$ alkyl,  $C_{1-4}$ alkyloxy, amino, hydroxy or halo; and

Het<sup>3</sup> [may also] represents furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl or imidazolyl each optionally substituted with one or two  $C_{1-4}$ alkyl substituents; pyrrolyl or pyrazolyl optionally substituted with formyl, hydroxy $C_{1-4}$ alkyl, hydroxycarbonyl,  $C_{1-4}$ alkyloxycarbonyl or with one or two  $C_{1-4}$ alkyl substituents; thiadiazolyl or oxadiazolyl optionally substituted with amino or  $C_{1-4}$ alkyl; pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl each optionally substituted with  $C_{1-4}$ alkyl,  $C_{1-4}$ alkyloxy, amino, hydroxy, halo, 4,5-dihydro-5-oxo-1H-tetrazolyl substituted

with C<sub>1-4</sub> alkyl, 2-oxo-3-oxazolidinyl, 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl or a radical of formula



wherein

A-Z represents S-CH=CH, S-CH<sub>2</sub>-CH<sub>2</sub>, S-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>, CH=CH-CH=CH, or CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>;


aryl is phenyl or phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, hydroxy, C<sub>1-4</sub> alkyl, polyhaloC<sub>1-4</sub> alkyl, cyano, aminocarbonyl, C<sub>1-4</sub> alkyloxy or polyhaloC<sub>1-4</sub> alkyloxy;

[provided] with the proviso that 5,6-dihydrospiro[imidazo[1,2-b][3]benzazepine-11[11H],4'-piperidine] and pharmaceutically acceptable addition salts thereof are not included.

4. (Amended) A compound according to [any one of the preceding] claim[s] 1 wherein -A-B- is a bivalent radical of formula -CH=CH-CH=CH- (a-3) or -CH=CH-Y- (a-2).
5. (Amended) A compound according to [any one of the preceding] claim[s] 1 wherein Z is - (CH<sub>2</sub>)<sub>p</sub>- (b-1), -CH=CH- (b-2), or -CH<sub>2</sub>-O- (b-4).
6. (Amended) A compound according to claim[s] 1, [2, 4 or 5] wherein L is hydrogen, C<sub>1-6</sub> alkyl, hydroxyC<sub>1-6</sub> alkyl, carboxyC<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkyloxycarbonyl, or C<sub>1-6</sub> alkyloxycarbonylC<sub>1-6</sub> alkyl.
7. (Amended) A compound according to [any one of the preceding] claim[s] 1 wherein R<sup>1</sup> is hydroxyC<sub>1-6</sub> alkyl, formyl, C<sub>1-6</sub> alkyloxycarbonyl, C<sub>1-6</sub> alkyloxyC<sub>1-6</sub> alkyl, N(R<sup>3</sup>R<sup>4</sup>)C(=O)-, halo or hydrogen.
8. (Amended) A compound according to claim 1 wherein the compound is 5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-carboxamide dihydrochloride;  
1'-butyl-5,6-dihydrospiro[imidazo[2,1-b][3] benzazepine-11-[11H],4'-piperidine];  
6,11-dihydro-1'-methylspiro[5H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine] cyclohexylsulfamate(1:2);

6,11-dihydrospiro[5-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-methanol] (E)-2-butenedioate (2:1);  
 3-chloro-6,11-dihydrospiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine] (E)-2-butenedioate (1:1);  
 6,11-dihydro-3-(methoxymethyl)spiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]. (E)-2-butenedioate (1:1);  
 6,11-dihydro-1'-(2-hydroxyethyl)spiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-carboxamide;  
 6,11-dihydro-1'-methylspiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-carboxamide monohydrate;  
 ethyl 3-(aminocarbonyl)-6,11-dihydro- $\alpha$ -phenylspiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-1'-propanoate monohydrochloride;  
 3-(aminocarbonyl)-6,11-dihydrospiro[5*H*-imidazo[2,1-b][3]-benzazepine-11,4'-piperidine]-1'-carboxylate;  
 spiro[10*H*-imidazo[1,2-a]thieno[3,2-d]azepine-10,4'-piperidine];  
 6,11-dihydrospiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-2,3-dicarboxamide dihydrochloride monohydrate; or  
 a prodrug, a *N*-oxide, an addition salt, a quaternary amine or a stereochemically isomeric form thereof.


10. (Amended) A pharmaceutical composition comprising a pharmaceutically acceptable carrier, and as active ingredient a therapeutically effective amount of a compound as [described] defined in [any one of] claim[s] 1 [to 9].
11. (Amended) A process of preparing a composition as claimed in claim 10, [characterized in that,] wherein a pharmaceutically acceptable carrier is [intimately] mixed with a therapeutically effective amount of a compound as [described] defined in [any one] claim[s] 1 [to 9].
12. (Amended) A compound of formula



or a *N*-oxide, an addition salt, a quaternary amine or a stereochemically isomeric form thereof wherein P is a protective group and n, -A-B-, Z, R<sup>1</sup> and R<sup>2</sup> are defined as in claim 1, [provided] with the proviso that 6,11-dihydro-1'-(phenylmethyl)-5*H*-spiro[imidazo[1,2-b][3]benzazepine-11,4'-piperidine] (E)-2-butenedioate(1:2) is not included.


14. (Amended) A process of preparing a compound as claimed in claim 1, [characterized by,] comprising

- a) deprotecting an intermediate of formula (II), followed optionally by derivatizing either the piperidine moiety, or the imidazole moiety, or both the piperidine moiety and the imidazole moiety




with [-A-B-, Z, L, R<sup>1</sup> and R<sup>2</sup>, and n defined as in claim 1 and] P being a protective group;

- b) derivatizing an intermediate of formula (II) at the imidazole moiety, [leading to the formation of] to form an intermediate of formula (II-a), followed by deprotecting the piperidine moiety, and followed optionally by derivatizing the piperidine moiety



[with -A-B-, Z, L, R<sup>1</sup> and R<sup>2</sup>, and n defined as in claim 1 and P being a protective group;]

- c) [by] cyclizing an intermediate of formula (III) in the presence of an appropriate acid,  
[resulting in] to form a compound of formula (I-a)




[with -A-B-, L, R<sup>1</sup> and R<sup>2</sup>, and n defined as in claim 1;]

and, [if desired] optionally, converting compounds of formula (I) and (I-a) into each other [following art-known transformations], and further, [if desired] optionally, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or [conversely,] converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid; and, [if desired] optionally, preparing stereochemically isomeric forms or N-oxide forms thereof.

15. (Amended) A process of preparing a compound as claimed in claim [12] 13, [characterized by] comprising,  
a) cyclizing a compound of formula (XI) with an appropriate acid, [leading] to form a compound of formula (II-k), followed optionally by derivatizing the imidazole moiety,




[leading] to form a compound of formula (II-a-1)



with [-A-B-, R<sup>1</sup>, R<sup>2</sup>, n and P defined as in claim 13, and] Z<sub>1</sub> being a bivalent radical of formula -(CH<sub>2</sub>)<sub>p</sub>-, wherein p is 1,2,3 or 4[.] ; and

- b) [by] reacting a tricyclic moiety of formula (XIV) with a reagent of formula (XV) under an inert atmosphere in a reaction inert solvent in the presence of a suitable base, [leading the] to form a compound of formula (II-l), followed optionally by derivatizing the imidazole moiety [leading] to form a compound of formula (II-a-2)



with [-A-B-, R<sup>1</sup>, R<sup>2</sup>, n and P defined as in claim 13,] W<sub>5</sub> being a suitable leaving group, [e.g. a halo,] and Z<sub>2</sub> being a bivalent radical of formula -(CH<sub>2</sub>)<sub>p</sub>-, or -CH<sub>2</sub>-O-, wherein p is 1,2,3 or 4.


# ANTI-HISTAMINIC SPIRO COMPOUNDS

The present invention is concerned with spiro compounds having antihistaminic activity. It further relates to their use as a medicine, their preparation as well as compositions comprising them.

WO 97/24356, published on 10 July 1997, discloses 4-(imidazo-azepine) piperidine spiro derivatives as intermediates in the preparation of 1-(1,2-disubstituted piperidiny)-4-(imidazo-azepine) piperidine spiro derivatives having tachykinin antagonistic activity.

Surprisingly, the 4-(imidazo-azepine) piperidine spiro derivatives of the present invention show an interesting antihistaminic activity profile.

The present invention concerns compounds of formula (I) for use as a medicine, characterized in that the compounds of formula (I) are defined as



their prodrugs, *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms wherein

$R^1$  is hydrogen,  $C_{1-6}$ alkyl, halo, formyl, carboxyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkyl-carbonyl,  $N(R^3R^4)C(=O)-$ ,  $N(R^3R^4)C(=O)N(R^5)-$ , ethenyl substituted with carboxyl or  $C_{1-6}$ alkyloxycarbonyl, or  $C_{1-6}$ alkyl substituted with hydroxy, carboxyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $N(R^3R^4)C(=O)-$ ,  $C_{1-6}alkylC(=O)N(R^5)-$ ,  $C_{1-6}alkylS(=O)_2N(R^5)-$  or  $N(R^3R^4)C(=O)N(R^5)-$ ;

wherein each  $R^3$  and each  $R^4$  independently are hydrogen or  $C_{1-4}$ alkyl;

$R^5$  is hydrogen or hydroxy;

$R^2$  is hydrogen,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy $C_{1-6}$ alkyl,  $N(R^3R^4)C(=O)-$ , aryl or halo;

$n$  is 1 or 2;

-A-B- represents a bivalent radical of formula



-2-

$$-\text{CH}=\text{CH}-\text{Y}- \quad (\text{a-2}); \text{ or}$$

$$-\text{CH}=\text{CH}-\text{CH}=\text{CH}- \quad (\text{a-3});$$

wherein each hydrogen atom in the radicals (a-1) to (a-3) may independently be

replaced by  $\text{R}^6$  wherein  $\text{R}^6$  is selected from  $\text{C}_{1-6}$ alkyl, halo, hydroxy,

5  $\text{C}_{1-6}$ alkyloxy, ethenyl substituted with carboxyl or  $\text{C}_{1-6}$ alkyloxycarbonyl, hydroxy $\text{C}_{1-6}$ alkyl, formyl, carboxyl and hydroxycarbonyl $\text{C}_{1-6}$ alkyl;

each Y independently is a bivalent radical of formula  $-\text{O}-$ ,  $-\text{S}-$  or  $-\text{NR}^7-$ ;

wherein  $\text{R}^7$  is hydrogen,  $\text{C}_{1-6}$ alkyl or  $\text{C}_{1-6}$ alkylcarbonyl;

Z is is a bivalent radical of formula

$$-(\text{CH}_2)_p- \quad (\text{b-1}),$$

$$-\text{CH}_2-\text{O}- \quad (\text{b-4}),$$

$$-\text{CH}=\text{CH}- \quad (\text{b-2}),$$

$$-\text{CH}_2-\text{C}(=\text{O})- \quad (\text{b-5}), \text{ or}$$

$$-\text{CH}_2-\text{CHOH}- \quad (\text{b-3}),$$

$$-\text{CH}_2-\text{C}(=\text{NOH})- \quad (\text{b-6}),$$

provided that the bivalent radicals (b-3), (b-4), (b-5) and (b-6) are connected to the nitrogen of the imidazole ring via their  $-\text{CH}_2-$  moiety;

wherein p is 1, 2, 3 or 4;

L is hydrogen;  $\text{C}_{1-6}$ alkyl;  $\text{C}_{2-6}$ alkenyl;  $\text{C}_{1-6}$ alkylcarbonyl;  $\text{C}_{1-6}$ alkyloxycarbonyl;

5  $\text{C}_{1-6}$ alkyl substituted with one or more substituents each independently selected from hydroxy, carboxyl,  $\text{C}_{1-6}$ alkyloxy,  $\text{C}_{1-6}$ alkyloxycarbonyl, aryl, aryloxy, cyano or  $\text{R}^8\text{HN}-$  wherein  $\text{R}^8$  is hydrogen,  $\text{C}_{1-6}$ alkyl,  $\text{C}_{1-6}$ alkyloxycarbonyl,  $\text{C}_{1-6}$ alkylcarbonyl; or

L represents a radical of formula

$$-\text{Alk}-\text{Y}-\text{Het}^1 \quad (\text{c-1}),$$

10  $-\text{Alk}-\text{NH}-\text{CO}-\text{Het}^2 \quad (\text{c-2}) \text{ or}$

$$-\text{Alk}-\text{Het}^3 \quad (\text{c-3}); \text{ wherein}$$

Alk represents  $\text{C}_{1-4}$ alkanediyl ;

Y represents O, S or NH ;

Het<sup>1</sup>, Het<sup>2</sup> and Het<sup>3</sup> each represent furanyl, tetrahydrofuranyl, thienyl, oxazolyl,

15 thiazolyl or imidazolyl each optionally substituted with one or two  $\text{C}_{1-4}$ alkyl substituents; pyrrolyl or pyrazolyl optionally substituted with formyl, hydroxy $\text{C}_{1-4}$ alkyl, hydroxycarbonyl,  $\text{C}_{1-4}$ alkyloxycarbonyl or with one or two  $\text{C}_{1-4}$ alkyl substituents;

thiadiazolyl or oxadiazolyl optionally substituted with amino or  $\text{C}_{1-4}$ alkyl; pyridinyl,


pyrimidinyl, pyrazinyl or pyridazinyl each optionally substituted with  $\text{C}_{1-4}$ alkyl,

20  $\text{C}_{1-4}$ alkyloxy, amino, hydroxy or halo; and

Het<sup>3</sup> may also represent 4,5-dihydro-5-oxo-1H-tetrazolyl substituted with  $\text{C}_{1-4}$ alkyl,

2-oxo-3-oxazolidinyl, 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl or a radical of formula

-3-



wherein

A-Z represents S-CH=CH, S-CH<sub>2</sub>-CH<sub>2</sub>, S-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>, CH=CH-CH=CH, or CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>;

aryl is phenyl or phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, hydroxy, C<sub>1-4</sub>alkyl, polyhaloC<sub>1-4</sub>alkyl, cyano, aminocarbonyl, C<sub>1-4</sub>alkyloxy or polyhaloC<sub>1-4</sub>alkyloxy.

The compounds of formula (I) are deemed novel provided that

5,6-dihydrospiro[imidazo[1,2-b][3]benzazepine-11[11H],4'-piperidine] and pharmaceutically acceptable addition salts thereof are not included and thus the present invention also relates to the compounds of formula (I) as defined hereinabove provided that 5,6-dihydrospiro[imidazo[1,2-b][3]benzazepine-11[11H],4'-piperidine] and pharmaceutically acceptable addition salts thereof are not included .

The term prodrug as used throughout this text means the pharmacologically acceptable derivatives, e.g. esters and amides, such that the resulting biotransformation product of the derivative is the active drug as defined in the compounds of formula (I). The reference by Goodman and Gilman (The Pharmacological Basis of Therapeutics, 8<sup>th</sup> ed., McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs", p. 13-15) describing prodrugs generally, is hereby incorporated.

As used herein C<sub>1-4</sub>alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as methyl, ethyl, propyl, 1-methylethyl, butyl and the like; C<sub>1-6</sub>alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the groups defined for C<sub>1-4</sub>alkyl and pentyl, hexyl, 2-methylpropyl, 2-methylbutyl and the like; C<sub>2-6</sub>alkenyl as a group or part of a group defines straight and branched chain hydrocarbon radicals containing one double bond and having from 2 to 6 carbon atoms such as ethenyl, 2-propenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-hexenyl and the like.

As used herein before, the term (=O) forms a carbonyl moiety when attached to a carbon atom and a sulfonyl moiety when attached twice to a sulfur atom. The term (=NOH) forms a hydroxylimine moiety when attached to a carbon atom.

The term halo is generic to fluoro, chloro, bromo and iodo. As used in the foregoing and hereinafter, polyhaloC<sub>1-4</sub>alkyl as a group or part of a group is defined as mono- or polyhalosubstituted C<sub>1-4</sub>alkyl, in particular methyl with one or more fluoro atoms, for example, difluoromethyl or trifluoromethyl. In case more than one halogen atoms are  
5 attached to an alkyl group within the definition of polyhaloC<sub>1-4</sub>alkyl, they may be the same or different.

When any variable (e.g. R<sup>3</sup>, R<sup>4</sup> etc.) occurs more than one time in any constituent, each  
10 definition is independent.

It will be appreciated that some of the compounds of formula (I), their prodrugs, N-oxides, addition salts, quaternary amines and stereochemically isomeric forms may contain one or more centers of chirality and exist as stereochemically isomeric forms.

15 The term "stereochemically isomeric forms" as used hereinbefore defines all the possible stereoisomeric forms which the compounds of formula (I), their prodrugs, N-oxides, addition salts, quaternary amines or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said  
20 mixtures containing all diastereomers and enantiomers of the basic molecular structure as well as each of the individual isomeric forms of formula (I), their prodrugs, N-oxides, addition salts or quaternary amines substantially free, *i.e.* associated with less than 10%, preferably less than 5%, in particular less than 2% and most preferably less than 1% of the other isomers. Stereochemically isomeric forms of the compounds of  
25 formula (I) are obviously intended to be embraced within the scope of this invention.

For therapeutic use, the addition salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases which are not-pharmaceutically acceptable may also find use, for example, in the  
30 preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

The pharmaceutically acceptable acid and base addition salts as mentioned hereinabove  
35 are meant to comprise the therapeutically active non-toxic acid and base addition salt forms which the compounds of formula (I) are able to form. The pharmaceutically acceptable acid addition salts can conveniently be obtained by treating the base form with such appropriate acid. Appropriate acids comprise, for example, inorganic acids

such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid, sulfuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic (i.e. ethanedioic), malonic, succinic (i.e. butanedioic acid), maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

Conversely said salt forms can be converted by treatment with an appropriate base into the free base form.

The compounds of formula (I) containing an acidic proton may also be converted into their non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

An interesting group of compounds consists of those compounds of formula (I) wherein L is hydrogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyloxycarbonyl or C<sub>1-6</sub>alkyl substituted with hydroxy, carboxyl, C<sub>1-6</sub>alkyloxy or C<sub>1-6</sub>alkyloxycarbonyl;

Another interesting group of compounds consists of those compounds of formula (I) wherein -A-B- is a bivalent radical of formula -CH=CH-CH=CH- (a-3).

Also interesting compounds are those compounds of formula (I) wherein -A-B- is a bivalent radical of formula -CH=CH-Y- (a-2).

Further interesting compounds are those compounds of formula (I) wherein Z is -(CH<sub>2</sub>)<sub>p</sub>- (b-1), -CH=CH- (b-2), or -CH<sub>2</sub>-O- (b-4).

Other interesting compounds are those compounds of formula (I) wherein L is hydrogen, C<sub>1-6</sub>alkyl, carboxyC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, or C<sub>1-6</sub>alkyloxycarbonylC<sub>1-6</sub>alkyl.

- 5 Again other interesting compounds are those compounds of formula (I) wherein L is hydroxyC<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyl substituted with aryl and C<sub>1-6</sub>alkyloxycarbonyl.

Also further interesting compounds are those compounds of formula (I) wherein R<sup>1</sup> is hydroxyC<sub>1-6</sub>alkyl, formyl, C<sub>1-6</sub>alkyloxycarbonyl, N(R<sup>3</sup>R<sup>4</sup>)C(=O)-, halo or hydrogen.

10

Other interesting compounds are those compounds of formula (I) wherein R<sup>1</sup> is C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl.

Special compounds are those compounds of formula (I) wherein one or more of the following restrictions apply :

15

-A-B- is a bivalent radical of formula -CH=CH-CH=CH- (a-3) wherein each hydrogen may independently be replaced by C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, halo or hydroxy;

Z is -(CH<sub>2</sub>)<sub>p</sub>- wherein p is 1,2,3 or 4, -CH<sub>2</sub>-C(=O)-, -CH<sub>2</sub>-CHOH-, -CH=CH-, -CH<sub>2</sub>-O-;

L is hydrogen, C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxycarbonyl;

20

R<sup>1</sup> is hydrogen, formyl, carboxyl, amide, halo, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, -NH-C(=O)-C<sub>1-6</sub>alkyl, -NH-C(=O)-NH<sub>2</sub>, -NH-SO<sub>2</sub>-C<sub>1-6</sub>alkyl;

R<sup>2</sup> is hydroxyC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, halo, amide.

25

The most preferred compounds are:

5,6-dihydrospiro[11*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-carboxamide dihydrochloride (comp. 17);

1'-butyl-5,6-dihydrospiro[imidazo[2,1-b][3] benzazepine-11-[11*H*],4'-piperidine] (comp. 3);

30

6,11-dihydro-1'-methylspiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine] cyclohexylsulfamate(1:2) (comp. 1);

6,11-dihydrospiro[5-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-methanol] (E)-2-butenedioate (2:1) (comp. 18a);


3-chloro-6,11-dihydrospiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]

35

(E)-2-butenedioate (1:1) (comp. 20);

6,11-dihydro-3-(methoxymethyl)spiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]. (E)-2-butenedioate (1:1) (comp. 58);

- 6,11-dihydro-1'-(2-hydroxyethyl)spiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-carboxamide (comp. 62);  
 6,11-dihydro-1'-methylspiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-carboxamide monohydrate (comp. 80);  
 5 ethyl 3-(aminocarbonyl)-6,11-dihydro- $\alpha$ -phenylspiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-propanoate monohydrochloride (comp. 64);  
 3-(aminocarbonyl)-6,11-dihydrospiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-carboxylate (comp. 79);  
 spiro[10*H*-imidazo[1,2-*a*]thieno[3,2-*d*]azepine-10,4'-piperidine] (comp. 56a);  
 10 6,11-dihydrospiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-2,3-dicarboxamide dihydrochloride monohydrate (comp. 53);  
 a prodrug, a *N*-oxide, an addition salt, a quaternary amine or a stereochemically isomeric form thereof.
- 15 The present invention also concerns novel compounds of formula



- their *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms, wherein
- $R^1$  is hydrogen,  $C_{1-6}$ alkyl, halo, formyl, carboxyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkyl-carbonyl,  $N(R^3R^4)C(=O)-$ ,  $N(R^3R^4)C(=O)N(R^5)-$ , ethenyl substituted with carboxyl or  $C_{1-6}$ alkyloxycarbonyl, or  $C_{1-6}$ alkyl substituted with hydroxy, carboxyl, amino,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $N(R^3R^4)C(=O)-$ ,  $C_{1-6}$ alkyl $C(=O)N(R^5)-$ ,  $C_{1-6}$ alkyl $S(=O)_2N(R^5)-$  or  $N(R^3R^4)C(=O)N(R^5)-$ ;  
 wherein each  $R^3$  and each  $R^4$  independently are hydrogen or  $C_{1-4}$ alkyl;  
 25  $R^5$  is hydrogen or hydroxy;  
 -A-B-, Z,  $R^2$ , and n are as defined for compounds of formula (I); and  
 P represents a protective group for example, benzyl, or those protective groups mentioned in Chapter 7 of "Protective Groups in Organic Synthesis" by T. Greene and P. Wuyts (John Wiley & Sons, Inc. 1991),  
 30 provided that 6,11-dihydro-1'-(phenylmethyl)-5*H*-spiro[imidazo[1,2-*b*][3]benzazepine-11,4'-piperidine] (E)-2-butenedioate(1:2) is not included.



The compounds of formula (II-a) are useful for the preparation of the compounds of formula (I).

Interesting compounds are those compounds of formula (II-a) wherein P is benzyl.

Also interesting compounds are those compounds of formula (II-a) wherein R<sup>1</sup> is hydrogen, halo, formyl, N(R<sup>3</sup>R<sup>4</sup>)C(=O)-, or C<sub>1-6</sub>alkyl substituted with hydroxy, amino, C<sub>1-6</sub>alkylC(=O)N(R<sup>5</sup>)-, C<sub>1-6</sub>alkylS(=O)<sub>2</sub>N(R<sup>5</sup>)- or N(R<sup>3</sup>R<sup>4</sup>)C(=O)N(R<sup>5</sup>)-.


Further interesting compounds are those compounds of formula (II-a) wherein -A-B- is a bivalent radical of formula (a-3) wherein each hydrogen atom may independently be replaced by C<sub>1-6</sub>alkyl, halo, hydroxy, or C<sub>1-6</sub>alkyloxy.

Again further interesting compounds are those compounds of formula (II-a) wherein Z is a bivalent radical of formula (b-1) and n is 1.


Compounds of formula (I), can be prepared by deprotecting an intermediate of formula (II), wherein P is a protecting group, for example, benzyl, or those protective groups mentioned in Chapter 7 of "Protective Groups in Organic Synthesis" by T. Greene and P. Wuyts (John Wiley & Sons, Inc. 1991). Said deprotection reaction can be performed by, for example, catalytic hydrogenation in the presence of hydrogen and an appropriate catalyst in a reaction inert solvent. A suitable catalyst in the above reaction is, for example, platinum-on-charcoal, palladium-on-charcoal, and the like. An appropriate reaction-inert solvent for said reaction is, for example, an alcohol, e.g. methanol, ethanol, 2-propanol and the like, an ester, e.g. ethylacetate and the like, an acid, e.g. acetic acid and the like. The thus obtained deprotected compounds can optionally further be derivatized either by replacing the hydrogen on the piperidine nitrogen by a moiety belonging to L, or by introducing on the imidazole moiety a R<sup>1</sup> group or a R<sup>2</sup> group or a R<sup>1</sup> and R<sup>2</sup> group, or by derivatizing both the piperidine moiety and the imidazole moiety.

An intermediate of formula II can also first be derivatized at the imidazole moiety by introducing a R<sup>1</sup> group or a R<sup>2</sup> group or a R<sup>1</sup> and R<sup>2</sup> group, resulting in an intermediate of formula (II-a), and then be deprotected, followed optionally by a derivation at the piperidine nitrogen.

-9-



Alternatively, compounds of formula (I), wherein Z is a bivalent radical of formula  $-\text{CH}_2-\text{C}(=\text{O})-$  (b-5), said compounds being represented by formula (I-a), can be prepared by cyclizing an intermediate of formula (III) in the presence of an acid, e.g. trifluoromethanesulfonic acid and the like.




The compounds of formula (I) may further be prepared by converting compounds of formula (I) into each other according to art-known group transformation reactions.

- 10 The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth
- 15 alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboxy-peroxy acid or halo substituted benzenecarboxy-peroxy acid, e.g. 3-chlorobenzenecarboxy-peroxy acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g.

t.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.


- 5 The compounds of formula (I) wherein L is other than hydrogen, said L being represented by  $L_a$  and said compounds being represented by formula (I-b) can be prepared by reacting the compounds of formula (I) wherein L is hydrogen, said compounds being represented by formula (I-c), with a reagent of formula  $L_a-W_1$  (IV), wherein  $W_1$  is a suitable leaving group, such as a halo atom, e.g. chloro, or
- 10 mesylate, tosylate, trifluoromethanesulfonate.



- Said reaction can conveniently be conducted in a reaction-inert solvent such as, for example, an aromatic hydrocarbon, an alkanol, a ketone, an ether, a dipolar aprotic solvent, a halogenated hydrocarbon, or a mixture of such solvents. The addition of an
- 15 appropriate base such as, for example, an alkali or an earth alkaline metal carbonate, hydrogen carbonate, alkoxide, hydride, amide, hydroxide or oxide, or an organic base, such as, for example, an amine, may be utilized to pick up the acid which is liberated during the course of the reaction. In some instances the addition of an iodide salt, preferably an alkali metal iodide, is appropriate. Somewhat elevated temperatures and
- 20 stirring may enhance the rate of the reaction.


- The compounds of formula (I-b), wherein  $L_a$  is optionally substituted  $C_{1-6}$ alkyl, said  $L_a$  being represented by  $L_{a1}$  and said compounds by formula (I-b-1), can be converted into the compounds of formula (I-c) by dealkylating and subsequently carbonylating the
- 25 compounds of formula (I-b-1) with a reagent of formula (V), wherein  $W_2$  represents a suitable leaving group, such as a halo atom, e.g. chloro, and  $L_{a2}$  represents  $C_{1-6}$ alkyloxycarbonyl, resulting in compounds of formula (I-b-2) and subsequently hydrolyzing the thus obtained compounds of formula (I-b-2).

-11-



- The reaction with reagent (V) is conveniently conducted by stirring and heating the starting material with the reagent in an appropriate solvent and in the presence of a suitable base. Appropriate solvents are, for example, aromatic hydro-carbons, e.g. methylbenzene, dimethylbenzene, chlorobenzene; ethers, e.g. 1,2-dimethoxyethane; methylenechloride and the like solvents. Suitable bases are, for example, alkali or earth alkaline metal carbonates, hydrogen carbonates, hydroxides, or organic bases such as, *N,N*-diethylethanamine, *N*-(1-methylethyl)-2-propanamine, and the like.
- The compounds of formula (I-b-2) are hydrolyzed in acidic or basic media following conventional methods. For example, concentrated acids such as hydrobromic, hydrochloric acid or sulfuric acid can be used, or alternatively bases such as alkali metal or earth alkaline metal hydroxides in water, an alkanol or a mixture of water-alkanol may be used. Suitable alkanols are methanol, ethanol, 2-propanol and the like.
- In order to enhance the rate of the reaction it is advantageous to heat the reaction mixture, in particular up to the reflux temperature.

- The compounds of formula (I-b-2) can also be prepared by reacting compounds of formula (I-c) with a reagent of formula (V) in the presence of a suitable base, e.g. *N,N*-diethylethanamine, in a reaction inert solvent, e.g. methylenechloride, or by reacting compounds of formula (I-c) with a reagent of formula (VI), e.g. *t*-butoxyanhydride, in a suitable solvent, such as, e.g. methylenechloride.



- The compounds of formula (I-b) wherein  $\text{R}^1$  or  $\text{R}^1$  and  $\text{R}^2$  represent hydroxymethyl, said compounds being represented by formula (I-b-3) and (I-b-4), can be prepared by


-12-

reacting the compounds of formula (I) wherein L is L<sub>a</sub> and R<sup>1</sup> and R<sup>2</sup> are hydrogen, said compounds being represented by formula (I-b-5), with formaldehyde, optionally in the presence of an appropriate carboxylic acid - carboxylate mixture such as, for example, acetic acid - sodium acetate and the like. In order to enhance the rate of the reaction, it is advantageous to heat the reaction mixture up to the reflux temperature.

The thus obtained compounds of formula (I-b-3) and (I-b-4) can be further oxidized to the corresponding aldehyde, represented by formula (I-b-6) and (I-b-7) or the corresponding carboxylic acid, represented by formula (I-b-8) and (I-b-9), by reaction with a suitable reagent such as, for example, manganese(IV)oxide, respectively, silver nitrate.


The compounds of formula (I-b-8) and (I-b-9) can further be converted in the corresponding amide, said compounds being represented by formula (I-b-10) and (I-b-11), by reaction with a suitable carbodiimide, e.g. 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, in the presence of ammonia and a suitable catalyst, e.g. *N,N*-dimethylaminopyridine, in a reaction inert solvent, e.g. methylenechloride.

-13-



The compounds of formula (I) wherein L is  $L_a$ , and  $R^1$  or  $R^1$  and  $R^2$  are halo, said compounds being represented by formula (I-b-12) and (I-b-13), can be prepared by


halogenating a compound of formula (I-b-5) with an appropriate halogenating reagent in a reaction-inert solvent.



A suitable halogenating reagent in the above reaction is, for example, an *N*-halogenated amide, e.g. *N*-bromosuccinimide. A suitable reaction-inert solvent for said halogenation reaction is, for example, *N,N*-dimethylformamide, *N,N*-dimethylacetamide, methylenechloride and the like. Another suitable halogenating reagent is, for example, tetrabutylammoniumtribromide in the presence of a suitable base, e.g. sodium carbonate, in a suitable solvent, e.g. 3-methyl-2-butanone or dichloromethane/water mixture.

Compounds of formula (I-b-12) can be converted in a compound of formula (I-b-10) by reaction with CuCN in the presence of a suitable solvent, e.g. *N,N*-dimethylformamide/water mixture.


The compounds of formula (I-b) wherein  $R^1$  is  $C_{1-6}$ alkyloxycarbonyl, said compounds being represented by formula (I-b-14), can be prepared by reacting a compound of formula (I-b-6) with a reagent of formula (VII) in the presence of a base e.g. piperidine, pyridine, and the like.




The compounds of formula (I-b-14) can further be hydrolyzed into a compound of formula (I-b) wherein  $R^1$  is carboxyethenyl, in the presence of an acid or a base in case  $L_a$  is  $C_{1-6}alkyl$ , or in the presence of a base, in case  $L_a$  is  $C_{1-6}alkyloxycarbonyl$ .

Compounds of formula (I-b) wherein  $L_a$  is substituted ethyl, said  $L_a$  being represented


by  $L_{a3}-CH_2-CH_2-$ , and said compounds by formula (I-b-15) can be prepared by reacting a compound of formula (I-c) with a reagent of formula (VIII) in the presence of a suitable base, e.g. sodium bicarbonate or triethylamine, in a suitable reaction inert solvent, e.g. *N,N*-dimethylformamide or a suitable alkanol, e.g. methanol.



Compounds of formula (I-c) may also be converted into compounds of formula (I-b-15) wherein  $L_{a3}$  is hydroxy, said compounds being represented by formula (I-b-15-1) by reaction with ethylene oxide in the presence of a suitable base, e.g. triethylamine, in a suitable reaction inert solvent, such as an alkanol, e.g. methanol.




Compounds of formula (I) wherein Z is a bivalent radical of formula  $-CH_2-CHOH-$  (b-3), said compounds being represented by formula (I-d), can be prepared by reducing a compound of formula (I-a) in the presence of a reducing reagent, e.g. sodium borohydride, in a reaction-inert solvent, e.g. methanol and the like.




Compounds of formula (I) wherein Z is a bivalent radical of formula  $-CH_2-C(=N-OH)-$  (b-6), said compounds being represented by the formula (I-e), can be prepared by



reacting a compound of formula (I-a) with hydroxylamine or a salt thereof, e.g. the hydrochloride salt, in a reaction inert solvent, e.g. pyridine and the like.



Compounds of formula (I) wherein Z is a bivalent radical of the formula  $-\text{CH}=\text{CH}-$  (b-2), said compounds being represented by formula (I-f), can be prepared by reacting a compound of formula (I-d) in the presence of an acid, e.g. methanesulfonic acid and the like.




Compounds of formula (I-b) wherein  $\text{L}_a$  is  $\text{C}_{1-6}$ alkyl substituted with  $\text{C}_{1-6}$ alkyloxy-carbonyl or  $\text{C}_{1-6}$ alkyloxycarbonylNH-, can be converted into compounds of formula (I-b) wherein  $\text{L}_a$  is  $\text{C}_{1-6}$ alkyl or amino $\text{C}_{1-6}$ alkyl by the hydrolysis reaction described above for the preparation of compounds of formula (I-c).

In the foregoing and the following preparations, the reaction mixture is worked up following art-known methods and the reaction product is isolated and, if necessary, further purified.


As described hereinabove, the intermediates of formula (II) can be derivatized at the imidazole moiety, said compounds being represented by formula (II-a), before being deprotected.

Introducing  $\text{R}^1$  or  $\text{R}^1$  and  $\text{R}^2$  wherein  $\text{R}^1$  and  $\text{R}^2$  represent hydroxymethyl, formyl, carboxyl or amide, in a compound of formula (II) can be performed as described hereinabove for the preparation of a compound of formula (I-b-3), (I-b-4), (I-b-6), (I-b-7), (I-b-8), (I-b-9), (I-b-10), (I-b-11).


- Intermediates of formula (II-a), wherein  $R^1$  is aminomethyl and  $R^2$  is hydrogen, said intermediates being represented by formula (II-b), can be prepared by reacting an intermediate of formula (II-c) with hydrogen and a mixture of methanol/ammonia in the presence of a suitable catalyst, for example rhodium on aluminium, in the presence of a catalyst poison, for example a thiophene solution.




- Intermediates of formula (II-a), wherein  $R^1$  is  $-\text{CH}_2\text{NHC}(=\text{O})\text{NH}_2$  and  $R^2$  is hydrogen, said intermediates being represented by formula (II-d) can be prepared by reacting an intermediate of formula (II-b) with potassium isocyanate in an appropriate acid, such as hydrochloric acid.




- Intermediates of formula (II-b) can also be converted in an intermediate of formula (II-a), wherein  $R^1$  is  $-\text{CH}_2\text{NHC}(=\text{O})\text{C}_{1-6}\text{alkyl}$  and  $R^2$  is hydrogen, said intermediate being represented by formula (II-e), by reaction with a reagent of formula (IX), wherein  $W_3$  represents a suitable leaving group, such as a halo atom, for example chloro, in the presence of a suitable base, e.g. *N,N*-diethylethanamine, in a reaction inert solvent, such as, for example methylenechloride.



Intermediates of formula (II-b) can further be converted into an intermediate of formula (II-a), wherein  $R^1$  is  $-\text{CH}_2\text{NHS}(=\text{O})_2\text{C}_{1-6}\text{alkyl}$  and  $R^2$  is hydrogen, said intermediate being represented by formula (II-f), by reaction with a reagent of formula (X) wherein  $W_4$  represents a suitable leaving group, such as a halo atom, e.g. chloro, in the presence of a suitable base, for example *N,N*-diethylethanamine, in a reaction inert solvent, such as methylenechloride.



Intermediates of formula (II) can also be halogenated according to the procedure described for the preparation of the compounds of formula (I-b-12) and (I-b-13), resulting in an intermediate of formula (II-g) and (II-h).




Intermediates of formula (II-g) and (II-h) can be converted in an intermediate of formula (II-i) and (II-j) by reaction under an atmosphere of ammonia and carbonmonoxide at elevated temperatures in the presence of a suitable catalyst, e.g.

acetic acid, palladium salt, and a suitable ligand, e.g. 1,3-bis(diphenylphosphino)-propane, in a reaction inert solvent, e.g. tetrahydrofuran.


In the following paragraphs, there are described several methods of preparing the starting materials in the foregoing preparations.

Intermediates of formula (II), wherein Z is a bivalent radical of formula  $-(CH_2)_p-$  (b-1), said Z being represented by  $Z_1$ , and said intermediates being represented by formula (II-k), can be prepared by cyclizing an alcohol of formula (XI). The intermediates of formula (II-k) may optionally be derivatized at the imidazole moiety resulting in an intermediate of formula (II-a-1) according to the procedures described for preparing a compound of formula (I-b-3), (I-b-4), (I-b-6) to (I-b-11) and an intermediate of formula (II-b) to (II-j).




Said cyclization reaction is conveniently conducted by treating the intermediate of formula (XI) with an appropriate acid, thus yielding a reactive intermediate which cyclizes to an intermediate of formula (II-k). Appropriate acids are, for example, strong acids, e.g. methanesulfonic acid, trifluoroacetic acid, and in particular superacid systems, e.g. trifluoromethanesulfonic acid, or Lewis acids, such as  $AlCl_3$  or  $SnCl_4$ . Obviously, only those compounds of formula II wherein P is stable under the given reaction conditions can be prepared according to the above reaction procedure.

Intermediates of formula (XI) can be prepared by reacting an imidazole derivative of formula (XII) with a ketone of formula (XIII).




Said reaction is conveniently performed in a reaction inert solvent such as, for example tetrahydrofuran, in the presence of a suitable base such as lithium diisopropylamide and butyl lithium.


- Intermediates of formula (II), wherein Z represents a bivalent radical of formula  $-(CH_2)_p-$  (b-1), or  $-CH_2-O-$  (b-4), said Z being represented by  $Z_2$ , and said intermediates being represented by formula (II-1), can also be prepared by reacting a tricyclic moiety of formula (XIV) with a reagent of formula (XV), wherein  $W_5$  represents a suitable leaving group, e.g. a halo atom, such as chloro, under an inert atmosphere in a reaction inert solvent, such as tetrahydrofuran, in the presence of a suitable base such as, for example, lithium diisopropylamide and butyl lithium. The intermediates of formula (II-1) may optionally be derivatized at the imidazole moiety resulting in an intermediate of formula (II-a-2) according to the procedures described for preparing a compound of formula (I-b-3), (I-b-4), (I-b-6) to (I-b-11) and an intermediate of formula (II-b) to (II-j).



- Intermediates of formula (XIV) can be prepared by cyclizing an intermediate of formula (XVI), according to the procedure described for the preparation of intermediates of formula (II-k).




Intermediates of formula (XVI) can be prepared by reduction from the corresponding aldehydes, said intermediates being represented by formula (XVII).




- Said reduction can be conducted in a suitable solvent, such as, for example methanol, in the presence of a suitable reducing agent, such as sodium borohydride.

Intermediates of formula (XVI) can also be prepared by reacting an intermediate of formula (XVIII) with formol under pressure.




Alternatively, the tricyclic moieties of formula (XIV), wherein Z represents a bivalent radical of formula  $-\text{CH}_2-$ , said Z being represented by  $\text{Z}_3$ , and said tricyclic moieties being represented by formula (XIV-a), may also be prepared by first cyclizing an intermediate of formula (XVII) wherein  $\text{Z}_2$  represents  $\text{Z}_3$ , said intermediate being represented by formula (XVII-a), by treating said intermediate with an appropriate acid, e.g. trifluoroacetic acid, leading to an intermediate of formula (XIX), followed by reduction in the presence of a suitable reducing agent.




Said reduction reaction can be performed in the presence of hydrogen and an appropriate catalyst in a reaction inert solvent. A suitable catalyst in the above reaction is, for example, platinum-on-charcoal, palladium-on-charcoal, and the like. An appropriate reaction inert solvent for said reaction is, for example, an alcohol, e.g. methanol, ethanol, 2-propanol and the like, an ester, e.g. ethylacetate and the like, an acid, e.g. acetic acid and the like.

Intermediates of formula (III) can be prepared by reacting an intermediate of formula (XX) with a reagent of formula (XXI), wherein  $\text{W}_6$  represents a suitable leaving group, such as a halo atom, e.g. chloro, in the presence of a suitable base, e.g. sodium hydride, in a reaction inert solvent, e.g. *N,N*-dimethylformamide and the like.




Intermediates of formula (XX), wherein  $R^1$  and  $R^2$  are hydrogen, said intermediates being represented by formula (XX-a), can be prepared by debenzylating an intermediate of formula (XXII).




- 5 Said debenzylation reaction can be performed by, for example, catalytic hydrogenation in the presence of hydrogen and an appropriate catalyst in a reaction inert solvent. A suitable catalyst in the above reaction is, for example, platinum-on-charcoal, palladium-on-charcoal, and the like. An appropriate reaction inert solvent for said reaction is, for example, an alcohol, e.g. methanol, ethanol, 2-propanol and the like, an ester, e.g. ethylacetate and the like, an acid, e.g. acetic acid and the like.

Intermediates of formula (XXII) can be prepared by imidazole formation out of an intermediate of formula (XXIII) in an acid, such as hydrochloric acid.




- 15 Intermediates of formula (XXIII) can be prepared by reacting an intermediate of formula (XXIV) with 2,2-dimethoxyethylamine in a reaction inert solvent, such as, for example *N,N*-dimethylformamide and the like.




Intermediates of formula (XXIV) can be prepared by reacting an intermediate of

formula (XXV) with thionylchloride.



Intermediates of formula (XXV) can be prepared by substituting an intermediate of formula (XXVI) with benzylamine in the presence of a suitable base, e.g. *N,N*-diethylethanamine, in a reaction inert solvent, e.g. methylenechloride and the like.



Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques, e.g., counter-current distribution, liquid chromatography and the like.

The compounds of formula (I) as prepared in the hereinabove described processes are generally racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) which are sufficiently basic or acidic may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid, respectively chiral base. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali or acid. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.



A number of intermediates and starting materials are commercially available or are known compounds which may be prepared according to conventional reaction procedures generally known in the art. For example, the preparation of 1-(1-phenyl-ethyl)-1*H*-imidazole is described in WO 92/22551.

5

The compounds of formula (I), their prodrugs, *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms possess useful pharmacological properties. In particular they are active antihistaminic agents, which activity can be demonstrated by for instance the 'Histamine - induced Lethality in Guinea Pigs' test (Arch. Int. Pharmacodyn. Ther., 251, 39-51, 1981), 'Protection of Rats from Compound 48/80 - induced Lethality' test (Arch. Int. Pharmacodyn. Ther., 234, 164-176, 1978), and 'Ascaris Allergy in Dogs' test (Arch. Int. Pharmacodyn. Ther., 251, 39-51, 1981 and Drug Dev. Res., 8, 95-102, 1986).

10

15

Some of the intermediates of formula (II-a) also have interesting pharmacological properties.

20

The compounds of the present invention have a selective binding affinity for the H<sub>1</sub> receptor, more in particular, they have a very low affinity for the 5HT<sub>2A</sub> serotonin receptor and the 5HT<sub>2C</sub> serotonin receptor. This dissociation between the H<sub>1</sub> receptor binding affinity and the 5HT<sub>2C</sub> and 5HT<sub>2A</sub> receptor binding affinity renders it unlikely for the present compounds to cause appetite stimulation and inappropriate weight gain reported for some other H<sub>1</sub>-antagonists.

25

An important asset of the present compounds is their lack of sedating properties at therapeutic dose levels, a troublesome side effect associated with many antihistaminic and antiallergic compounds. The non-sedating properties of the present compounds can be demonstrated, for example, by the results obtained in studying the sleep - wakefulness cycle of the rat (Psychopharmacology, 97, 436-442, (1989)) and the state of vigilance using EEG power spectra in wake rats (Sleep Research 24A, 118, (1995)).

30

The compounds of the present invention are also characterized by the absence of relevant cardio-hemodynamic and electrophysiological effects such as QTc prolongation.

35

An additional advantage of some of the present compounds is that they exhibit little or no metabolic transformations in animal and human liver, thus indicating a low risk for metabolic interactions.

Another interesting feature of the present compounds relates to their fast onset of action and the favorable duration of their action. The latter characteristic may enable the administration of the compound once daily.

5

The present compounds have a favorable physicochemical profile, particularly in terms of solubility and chemical stability.

10

In view of their physicochemical and pharmacological properties, the compounds of formula (I), their prodrugs, *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms thereof are very useful in the treatment of a broad range of allergic diseases such as, for example, allergic rhinitis, allergic conjunctivitis, chronic urticaria, pruritis, allergic asthma and the like.

15

Also in view of their useful physicochemical and pharmacological properties the subject compounds may be formulated into various pharmaceutical forms for administration purposes. To prepare the antiallergic compositions of this invention, an effective amount of the particular compound, in base or acid addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are, desirably as unitary dosage forms, administered orally, parenterally, percutaneously, rectally or topically for systemic action, or for topical action. In case of oral liquid pharmaceutical preparations, comprising solutions, suspensions, syrups, elixirs and emulsions, any of the usual pharmaceutical media, such as, for example, water, glycols, oils, alcohols and the like, may be employed, whereas in case of oral solid pharmaceutical preparations, comprising powders, pills, capsules and tablets, excipients such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like may be employed. Because of their ease in administration, tablets and capsules represent the most advantageous oral unit dosage forms, in which case solid pharmaceutical carriers are obviously employed. In case of injectable pharmaceutical compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, such as semipolair solvents, may be included, for example, to aid solubility. Examples of carriers for injectable solutions comprise saline solution, glucose solution or a mixture of saline and glucose solution. Injectable solutions containing compounds of the aforementioned formulas may also be formulated in an oil for prolonged action. Appropriate oils for this purpose are, for example, peanut oil, sesame oil, cottonseed oil, corn oil, soy bean oil, synthetic glycerol esters of long chain

20

25

30

35

fatty acids and mixtures of these and other oils. For the preparation of injectable suspensions, appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wettable agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause any significant deleterious effects on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment or as a gel. In case of pharmaceutical compositions for rectal administration, any of the usual excipients may be employed, comprising fat based and water soluble excipients, optionally combined with suitable additives, such as suspending or wetting agents. As appropriate compositions for topical application there may be cited all compositions usually employed for topically administering drugs e.g. creams, gellies, dressings, lotions, shampoos, tinctures, pastes, ointments, salves, ovules, powders, inhalations, nose sprays, eye drops and the like. Semisolid compositions such as salves, creams, gellies, ointments and the like will conveniently be used, but application of said compositions may be, for example, also by aerosol, e.g. with a propellant such as nitrogen, carbon dioxide, a freon, or without a propellant such as a pump spray or drops.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, powder packets, suppositories, ovules, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

The present invention also relates to a method of treating warm-blooded animals suffering from allergic diseases by administering to said warm-blooded animals an effective anti-allergic amount of a compound of formula (I), a prodrug, *N*-oxide, addition salt, quaternary amine or stereochemically isomeric form thereof.

The present invention further relates to the compounds of formula (I), their prodrugs, *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms

thereof for use as a medicine, and hence, the use of the present compounds for the manufacture of a medicament for treating warm-blooded animals suffering from allergic diseases is also part of the present invention.

In general it is contemplated that an effective antiallergic amount would be from about 0.001 mg/kg to about 2 mg/kg body weight, and more preferably from about 0.01 mg/kg to about 0.5 mg/kg body weight. In any event, an effective antiallergic amount may depend on the type and severity of the affliction to be treated and the evaluation of the physician prescribing the treatment with the subject drugs.

The following examples are intended to illustrate the scope of the present invention.

#### Experimental part

Hereinafter, THF means tetrahydrofuran, DIPE means diisopropyl ether, DMF means *N,N*-dimethylformamide, DIPA means diisopropyl amine

#### A. Preparation of intermediate compounds

##### Example A1

a) A mixture of DIPA (1.4 mol) in THF (3000ml) was stirred at -70°C under N<sub>2</sub> flow. Butyllithium 2.5 M/hexane (1.3 mol) was added portionwise at a temperature below -40°C. The mixture was stirred at -70°C for 15 min. 1-phenylethyl-1*H*-imidazole (1 mol) dissolved in THF was added dropwise at a temperature below -55°C. The mixture was stirred at -70°C for 1 hour. 1-(phenylmethyl)-4-piperidinone (1.2 mol) dissolved in THF was added dropwise at a temperature below -55°C. The mixture was stirred at -70°C for 1 hour, then brought to room temperature, stirred at room temperature overnight and decomposed with H<sub>2</sub>O. The organic solvent was evaporated. The aqueous concentrate was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was crystallized from DIPE (1100ml). The precipitate was filtered off, washed with DIPE and dried, yielding 271g of 4-[1-(2-phenylethyl)-1*H*-imidazol-2-yl]-1-(phenylmethyl)-4-piperidinol (75%) (interm. 1).

b) A mixture of intermediate (1) (0.75 mol) in trifluoromethanesulfonic acid (1500ml) was stirred at 65°C for 120 hours, then cooled, poured out on ice, alkalized with NaOH 50% and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was crystallized from DIPE/CH<sub>3</sub>CN (99/1) (1200ml). The precipitate was filtered off and dried, yielding 169.6g of 5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine] (66%) (interm. 2).

Example A2

1-(phenylmethyl)-4-[1-(phenylmethyl)-1*H*-imidazol-2-yl]-4-piperidinol (0.124 mol) and AlCl<sub>3</sub> (0.31 mol) were stirred in a melt at 120°C for 1h. The mixture was cooled, AlCl<sub>3</sub> (0.31 mol) was added and the mixture was stirred at 120°C for 1h. The mixture  
5 was poured into ice, alkalized with NaOH 50% and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried (MgSO<sub>4</sub>), filtered off and evaporated. The residue was purified by HPLC (eluent : CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 99/1). The pure fractions were collected and evaporated. The residue was converted into the hydrochloric acid salt (1:2) in (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O, yielding 0.91g of 1'-(phenylmethyl)spiro[imidazo[1,2-*b*]isoquinoline-  
10 10[5*H*],4'-piperidine]dihydrochloride.dihydrate (2%) (interm. 3; mp. 161.2 °C).

Example A3

a) A mixture of intermediate (2) (0.09 mol) in CH<sub>2</sub>Cl<sub>2</sub> (1000ml) was cooled to 0°C. 1-bromo-2,5-pyrrolidinedione (0.09 mol) was added portionwise over a 1-hour period. The organic layer was separated, washed with H<sub>2</sub>O, dried, filtered and the solvent was  
15 evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 97.5/2.5 to 95/5). A pure fraction was collected and the solvent was evaporated. The residue was dissolved in ethanol and converted into the (E)-2-butenedioic acid salt (1:1). The precipitate was filtered off and dried, yielding 17.3g of 3-bromo-5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]-  
20 benzazepine-11,4'-piperidine (E)-2-butenedioate(1:1) (36%) (interm. 4). Part of this fraction (16.5g) was taken up in H<sub>2</sub>O, K<sub>2</sub>CO<sub>3</sub> and CH<sub>2</sub>Cl<sub>2</sub>. The mixture was separated into its layers. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layer was dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated, yielding 12.9g of 3-bromo-5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-  
25 piperidine (interm. 4a).  
b) A mixture of intermediate (4a) (0.21 mol), 1,3-propanediylbis[diphenylphosphine] (2.5 g) and acetic acid, palladium(2+) salt (0.68 g) in THF (567 ml) was stirred in an autoclave at 150°C for 16 hours under NH<sub>3</sub> (10 atm) and CO (30 atm). The mixture was filtered and the filtrate was evaporated. This residue was purified by column  
30 chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/C<sub>2</sub>H<sub>5</sub>OH 100/0 over 46 min to 70/30). The pure fractions were collected and the solvent was evaporated, yielding 36 g of 5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-carboxamide (44%) (interm. 5).

Example A4

35 a) A mixture of intermediate (2) (0.16 mol) and sodium acetate (45g) in formol 38% (300ml) and acetic acid (30ml) was stirred and refluxed for 6 hours, then cooled,

poured out on ice and alkalized with a NaOH solution. The precipitate was filtered off and purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 99/1). The desired fractions were collected and the solvent was evaporated. The residue was triturated in CH<sub>3</sub>CN, filtered off and dried, yielding 13g of 5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-methanol (interm. 6).

b) A mixture of intermediate (6) (0.032 mol) and MnO<sub>2</sub> (65g) in chloroform (250ml) was stirred and refluxed for 2 hours, then cooled, filtered over dicalite and the filtrate was evaporated, yielding 11g of 5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-carboxaldehyde (interm. 7).

c) A mixture of intermediate (7) (0.0296 mol) in CH<sub>3</sub>OH/NH<sub>3</sub> (500ml) was hydrogenated at 50°C with Rh/Al<sub>2</sub>O<sub>3</sub> 5% (2g) as a catalyst in the presence of a thiophene solution (2ml). After uptake of H<sub>2</sub> (1 equiv), the catalyst was filtered off and the filtrate was evaporated, yielding 11g of 5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-methanamine (interm. 8). Part of this fraction (1g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 95/5). The pure fractions were collected and the solvent was evaporated. The residue was dissolved in 2-propanol and converted into the hydrochloric acid salt (1:3) with 2-propanol/HCl. The mixture was crystallized from DIPE. The precipitate was filtered off and dried, yielding 0.8g of 5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-methanamine hydrochloride (1:3) hydrate (1:1) (interm. 8a).

d) A mixture of intermediate (8) (0.0198 mol) in HCl 1N (50ml) was stirred at 50°C. KOCN (0.023 mol) was added portionwise (4x0.5g). The mixture was stirred at 50°C for 2 hours, then cooled, neutralized with a NaHCO<sub>3</sub> solution and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, washed with H<sub>2</sub>O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 95/5). The pure fractions were collected and the solvent was evaporated, yielding 4.3g of *N*-[[5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepin]-3-yl]methyl]methyl]urea (interm. 9)

#### Example A5

A mixture of intermediate (8) (0.0295 mol) and triethylamine (0.035 mol) in CH<sub>2</sub>Cl<sub>2</sub> (140ml) was stirred at room temperature. A solution of acetyl chloride (0.03 mol) in CH<sub>2</sub>Cl<sub>2</sub> (10ml) was added dropwise. The mixture was stirred at room temperature for 1 hour and poured out into H<sub>2</sub>O. K<sub>2</sub>CO<sub>3</sub> (2g) was added. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, washed with H<sub>2</sub>O, dried, filtered and

the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 97/3). The pure fractions were collected and the solvent was evaporated. Part of the residue (1.3g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was triturated in DIPE, filtered off and dried, yielding *N*-[[5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]-benzazepine-11,4'-piperidin]-3-yl]methyl]acetamide (interm. 10).

#### Example A6

A mixture of intermediate (8) (0.012 mol) and triethyl amine (0.015 mol) in CH<sub>2</sub>Cl<sub>2</sub> (150ml) was stirred at 0°C under N<sub>2</sub> flow. Methanesulfonyl chloride (0.013 mol) was added dropwise. The mixture was stirred for 2 hours. H<sub>2</sub>O was added and the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, washed with H<sub>2</sub>O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 98/2). The pure fractions were collected and the solvent was evaporated, yielding 2.1g of *N*-[[5,6-dihydro-1'-(phenylmethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidin]-3-yl]methyl]methanesulfonamide (interm. 11).

#### Example A7

a) 1-Methyl-4-phenyl-4-piperidinecarbonyl chloride (0.49 mol) was added portionwise at room temperature to a stirring mixture of benzenemethanamine (0.49 mol) and triethyl amine (1.223 mol) in CH<sub>2</sub>Cl<sub>2</sub> (2500ml). The mixture was stirred at room temperature for 1 hour. K<sub>2</sub>CO<sub>3</sub> (150g) and H<sub>2</sub>O were added. The mixture was stirred and separated into its layers. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layer was dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated, yielding 144g of 1-methyl-4-phenyl-*N*-(phenylmethyl)-4-piperidinecarboxamide (95%) (interm. 12).

b) A mixture of intermediate (12) (0.47 mol) in thionylchloride (750ml) was stirred and refluxed for 1 hour. The solvent was evaporated. Toluene was added twice and evaporated again, yielding 190g of *N*-[chloro(1-methyl-4-phenyl-4-piperidinyl)-methylene]benzenemethanamine monohydrochloride (100%) (interm. 13).

c) A mixture of intermediate (13) (0.47 mol) in DMF (750ml) was cooled on an ice bath. 2,2-Dimethoxyethanamine (0.54 mol) dissolved in DMF was added dropwise. The mixture was stirred at room temperature overnight. The solvent was evaporated, yielding 210g of *N*-(2,2-dimethoxyethyl)-1-methyl-4-phenyl-*N'*-(phenylmethyl)-4-piperidinecarboximidamide dihydrochloride (100%) (interm. 14).

d) A mixture of intermediate (14) (0.47 mol) in HCl 6N (1500ml) was stirred until a cloudy solution, then washed with CH<sub>2</sub>Cl<sub>2</sub> (900ml), stirred at 80°C for 1 hour, cooled, alkalized with a NaOH 50% solution and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was crystallized from CH<sub>3</sub>CN. The precipitate was filtered off and dried, yielding 38.3g of 1-methyl-4-phenyl-4-[1-(phenylmethyl)-1*H*-imidazol-2-yl]piperidine (25%) (interm. 15).

e) A mixture of intermediate (15) (0.195 mol) in methanol (350ml) was hydrogenated at room temperature for 18 hours with palladium on charcoal 10% (3g) as a catalyst.

After uptake of H<sub>2</sub> (1 equiv), the catalyst was filtered off and the filtrate was evaporated. The residue was crystallized from CH<sub>3</sub>CN. The precipitate was filtered off and dried, yielding 42.3g of 4-(1*H*-imidazol-2-yl)-1-methyl-4-phenylpiperidine (90%) (interm. 16).

f) A mixture of sodium hydride 60% (0.232 mol) in DMF (150ml) was stirred at room temperature. Intermediate (16) (0.145 mol) dissolved in DMF (400ml) was added dropwise. The mixture was stirred at room temperature for 1 hour. Methyl 2-chloroacetate (0.232 mol) dissolved in DMF (400ml) was added dropwise. The mixture was stirred at room temperature for 20 min, poured out into a solution of NaHCO<sub>3</sub> (20g) in H<sub>2</sub>O (2000ml) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 95/5). The pure fractions were collected and the solvent was evaporated, yielding 40.1g of methyl 2-(1-methyl-4-phenyl-4-piperidinyl)-1*H*-imidazole-1-acetate (88%) (interm. 17).

#### Example A8

a) Reaction under N<sub>2</sub> atmosphere. A mixture of DIPA (0.455 mol) in THF (500 ml) was stirred at -78°C. Butyllithium, 2.5M/hexane (0.390 mol) was added dropwise at -40°C. The mixture was stirred for 15 min, then re-cooled to -78°C. A solution of 1-(4-phenylbutyl)-1*H*-imidazole, prepared according to the procedure described in J. Chem. Soc., Perkin Trans., 1 (1975), 17, 1670-1671, (0.325 mol) in THF (350 ml) was added dropwise at -60°C. The mixture was stirred for one hour, then re-cooled to -78°C. This mixture was added dropwise to a mixture of *N,N*-dimethylformamide (0.390 mol, dry, p.a.) in THF (500 ml), stirred at -78°C. The reaction mixture was stirred for one hour at -78°C, then allowed to warm to room temperature while stirring overnight. A saturated aqueous NH<sub>4</sub>Cl solution (400 ml) was added and this mixture was extracted with THF. The separated organic layer was dried, filtered and the solvent



evaporated, yielding 74.2 g of 1-(4-phenylbutyl)-1*H*-imidazole-2-carboxamide (interm. 18).

b) A mixture of intermediate (18) (0.325 mol) in methanol (1400 ml) was stirred at room temperature. NaBH<sub>4</sub> (0.650 mol) was added portionwise and the reaction mixture was stirred for 2 hours at room temperature. The solvent was evaporated. The residue was taken up into water and this mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The separated organic layer was dried, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 99/1, 97/3, 96/4, 95/5 and 93/7). The desired fractions were collected and the solvent was evaporated, yielding 49.5 g of 1-(4-phenylbutyl)-1*H*-imidazole-2-methanol (66%) (interm. 19).

c) A mixture of intermediate (19) (0.417 mol) in methanesulfonic acid (960ml) was stirred at 120°C for 40 hours, then cooled, poured out on ice and alkalinized with NH<sub>4</sub>OH. The organic layer was separated, dried, filtered and the solvent evaporated. This fraction was purified by HPLC over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 97/3). A pure fraction was collected and the solvent was evaporated. The residue was crystallized from DIPE. The precipitate was filtered off and dried, yielding 15.7g of 6,7,8,13-tetrahydro-5*H*-imidazo[2,1-*b*][3]benzazonine (18%) (interm. 20).

d) A mixture of DIPA (0.151 mol) in THF (650ml) was stirred at -78°C under N<sub>2</sub> flow. Butyllithium 2,5M in hexane (0.144 mol) was added dropwise at a temperature below -40°C. The mixture was stirred at -78°C for 15 min. Intermediate (20) (0.072 mol) in a small amount of THF was added dropwise at a temperature below -55°C. The mixture was stirred at -78°C for 1 hour. *N,N*-bis(2-chloroethyl)benzenemethanamine hydrochloride in a small amount of THF was added dropwise at a temperature below -50°C. The mixture was stirred at -78°C for 1 hour, allowed to warm to room temperature overnight and decomposed with H<sub>2</sub>O. The organic solvent was evaporated. The aqueous concentrate was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried, filtered and the solvent was evaporated. This fraction was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 99/1, 98/2, 96/4, 94/6 and 92/8). A fraction was collected and the solvent was evaporated, yielding 5,6,7,8-tetrahydro-1'-(phenylmethyl)spiro[13*H*-imidazo[2,1-*b*][3]benzazonine-13,4'-piperidine] (interm. 21).

#### Example A9

A mixture of 5,10-dihydro-imidazo[1,2-*b*]isoquinoline-7,8-diol, obtained according to the procedure described in Ex.No. A8 c, (0.155 mol), phenyltrimethylammonium chloride (0.31 mol) and K<sub>2</sub>CO<sub>3</sub> (0.68 mol) in DMF (400ml) was stirred at 90°C for 20

hours, cooled, poured out into H<sub>2</sub>O and filtered over dicalite. The filtrate was separated into its layers. The organic layer was washed with H<sub>2</sub>O, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0 to 97/3). The pure fractions were collected and the solvent was evaporated, yielding 3g of 5,10-dihydro-7,8-dimethoxyimidazo[1,2-b]isoquinoline (8.4%) (interm. 22).

#### Example A10


A mixture of intermediate (31) (see Table 1), prepared according to the procedure described in Ex.No. A8d, (0.01 mol) in HBr 48 % solution (60ml) was stirred and refluxed for 2 hours. The solvent was evaporated. The residue was taken up in a small amount of H<sub>2</sub>O. The mixture was alkalized with K<sub>2</sub>CO<sub>3</sub> and extracted with CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH. The organic layer was separated, dried, filtered and the solvent was evaporated, yielding 4.3g of 5,6-dihydro-1'-(phenylmethyl)spiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-8,9-diol (100%) (interm. 23).

#### Example A11

A mixture of compound (22) (0.0117 mol) and triethyl amine (0.0421 mol) in toluene (100ml) was stirred and refluxed. Ethyl carbonochloridate (0.0702 mol) was added dropwise at reflux temperature. The mixture was stirred and refluxed for 1 hour, cooled, poured out into H<sub>2</sub>O and K<sub>2</sub>CO<sub>3</sub> (15g) and separated into its layers. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layer was dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/ethanol 96/4). The pure fractions were collected and the solvent was evaporated. The residue was boiled in DIPE. The precipitate was filtered off and dried, yielding 2.4g of [1'-(ethoxycarbonyl)spiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-6-yl] ethyl carbonate (33%) (interm. 24).

Table 1 lists intermediates which were prepared according to one of the above mentioned examples.

#### Table 1



Interm. No.	Ex. No.	n	z	R1	-A-B-
25	A8d	2	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-CH=CH-CH=CH-
3	A2	1	-CH <sub>2</sub> -	H	-CH=CH-CH=CH-
2	A1b	1	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-CH=CH-CH=CH-
26	A8d	1	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-CH=CF-CH=CH-
27	A8d	1	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-CH=CH-CH=CCH <sub>3</sub> -
28	A8d	1	-(CH <sub>2</sub> ) <sub>3</sub> -	H	-CH=CH-CH=CH-
29	A8d	1	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-COH=CH-CH=CH-
30	A8d	1	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-CH=CH-COH=CH-
31	A8d	1	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-CH=COCH <sub>3</sub> -COCH <sub>3</sub> =CH-
32	A8d	1	-O-CH <sub>2</sub> -	H	-CH=CH-CH=CH-
23	A10	1	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-CH=COH-COH=CH-
6	A4a	1	-(CH <sub>2</sub> ) <sub>2</sub> -	CH <sub>2</sub> OH	-CH=CH-CH=CH-
7	A4b	1	-(CH <sub>2</sub> ) <sub>2</sub> -	C(=O)H	-CH=CH-CH=CH-
8	A4c	1	-(CH <sub>2</sub> ) <sub>2</sub> -	CH <sub>2</sub> NH <sub>2</sub>	-CH=CH-CH=CH-
10	A5	1	-(CH <sub>2</sub> ) <sub>2</sub> -	CH <sub>2</sub> NHC(=O)CH <sub>3</sub>	-CH=CH-CH=CH-
9	A4d	1	-(CH <sub>2</sub> ) <sub>2</sub> -	CH <sub>2</sub> NHC(=O)NH <sub>2</sub>	-CH=CH-CH=CH-
5	A3b	1	-(CH <sub>2</sub> ) <sub>2</sub> -	C(=O)NH <sub>2</sub>	-CH=CH-CH=CH-
21	A8d	1	-(CH <sub>2</sub> ) <sub>4</sub> -	H	-CH=CH-CH=CH-
11	A6	1	-(CH <sub>2</sub> ) <sub>2</sub> -	CH <sub>2</sub> NHSO <sub>2</sub> CH <sub>3</sub>	-CH=CH-CH=CH-
33	A8d	1	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-CH=COCH <sub>3</sub> -COCH <sub>3</sub> =CH-

### B. Preparation of final compounds

#### Example B1 (preparative example)

- 5 A mixture of intermediate (2) (0.02 mol) in methanol (150ml) was hydrogenated with palladium on charcoal 10% (2g) as a catalyst at 50°C for 18hours. After uptake of H<sub>2</sub> (1eq), the catalyst was filtered and the filtrate was evaporated, yielding 5,6-dihydro-spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine] (comp. 6; not claimed). This fraction was converted into the hydrochloric acid salt (1:1) in CH<sub>3</sub>CN, yielding 5g of 5,6-dihydrospiro[imidazo[1,2-*b*][3]benzazepine-11[11*H*],4'-piperidine]
- 10 monohydrochloride (86%) (comp. 6a; not claimed). A fraction obtained in said way, can also be converted into the (E)-2-butenedioic acid salt.

#### Example B2

- a) A mixture of compound (6) (0.1 mol) and *N,N*-diethylethanamine (0.13 mol) in CH<sub>2</sub>Cl<sub>2</sub> (300ml) was stirred at a temperature below 10°C. Ethyl carbonochloridate
- 15 (0.12 mol) was added dropwise at this temperature. The mixture was allowed to warm

to room temperature and then stirred at room temperature for 1 hour. Water and  $K_2CO_3$  (10g) were added. The mixture was separated into its layers. The aqueous layer was extracted with  $CH_2Cl_2$ . The combined organic layer was dried ( $MgSO_4$ ), filtered and the solvent was evaporated, yielding 35.4g of ethyl 5,6-dihydrospiro[11*H*-imidazo-

5 [2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-carboxylate (100%) (comp. 4).

b) A mixture of compound (4) (0.1 mol), sodium acetate (0.3 mol) and acetic acid (0.258 mol) in formaldehyde 38% solution (165ml) was stirred and refluxed for 10 hours. The mixture was poured out into ice and a NaOH solution and extracted with  $CH_2Cl_2$ . The organic layer was separated, dried ( $MgSO_4$ ), filtered and the solvent was

10 evaporated. The residue was purified by column chromatography over silica gel (eluent:  $CH_2Cl_2$ /ethanol 95/5 to 90/10). The pure fractions were collected and the solvent was evaporated, yielding 16.5g of ethyl 5,6-dihydro-3-(hydroxymethyl)spiro-

[11*H*-imidazo[2,1-*b*][3]-benzazepine-11,4'-piperidine]-1'-carboxylate (46%) (comp. 5).

c) A mixture of compound (5) (0.046 mol) and potassium hydroxide (0.46 mol) in

15 2-propanol (130ml) was stirred and refluxed for 7 hours. The solvent was evaporated. The residue was taken up in water and extracted with  $CH_2Cl_2$ . The organic layer was separated, dried ( $MgSO_4$ ), filtered and the solvent was evaporated, yielding 11.5g of 5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-methanol (88%) (comp. 18). Part of this fraction (1g) was dissolved in  $CH_3OH$  and converted

20 into the (E)-2-butenedioic acid salt (2:1). The precipitate was filtered off and dried, yielding 0.6g of 5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-methanol (E)-2-butenedioic acid salt (2:1) (comp. 18a).

#### Example B3

A mixture of compound (6) (0.01 mol) and  $(CH_2O)_n$  (0.066 mol) in methanol (150ml)

25 and thiophene 4% solution (1ml) was hydrogenated with palladium on charcoal 10% (1g) as a catalyst at 50°C. After uptake of  $H_2$  (1eq), the catalyst was filtered and the filtrate was evaporated. The residue was taken up in  $H_2O/K_2CO_3/NH_4OH$  and stirred. The mixture was extracted with  $CH_2Cl_2$ , dried, filtered and evaporated. The residue was converted into the cyclohexanesulfamic acid salt (1:2) in 2-propanone and

30 recrystallized twice from 2-propanol, yielding 2.44g of 6,11-dihydro-1'-methyl-spiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine] cyclohexylsulfamate(1:2) (40%) (comp. 1).

#### Example B4

a) A mixture of 1-bromobutane (0.012 mol), compound (6) (0.01 mol),  $Na_2CO_3$

35 (0.02 mol) and potassium iodide (few crystals) in 2-butanone (200ml) was stirred and refluxed overnight. The mixture was evaporated, the residue was taken up in water and

extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was dried, filtered off and evaporated. The residue was purified by column chromatography over silica gel (eluent :  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  95/5). The pure fractions were collected and evaporated. The residue was converted into the hydrochloric acid salt (1:2) in 2-propanol. The precipitate was filtered and dried, yielding 0.7g of 1'-butyl-5,6-dihydrospiro[imidazo[2,1-b][3]-benzazepine-11-[11H],4'-piperidine] dihydrochloride.hemihydrate (18%) (comp. 3).

b) A mixture of 1-(3-chloropropoxy)-4-fluorobenzene (0.018 mol), 5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine] (0.015 mol),  $\text{Na}_2\text{CO}_3$  (0.015 mol) and KI (10 mg) in 4-methyl-2-pentanone (200ml) was stirred and refluxed for 18 hours. The mixture was poured into water, separated and the aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was dried ( $\text{MgSO}_4$ ), filtered off and evaporated. The residue was purified on a glass filter over silica gel (eluent :  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  90/10). The pure fractions were collected and evaporated. The residue was converted into the cyclohexylsulfamic acid salt (1:2) in 2-propanone. Yielding : 4.26g of 1'-[3-(4-fluorophenoxy)propyl]-5,6-dihydrospiro[imidazo[2,1-b][3]benzazepine-11-[11H],4'-piperidine] cyclohexylsulfamate(1:2) (37%); mp. 180°C (comp. 71).

c) A mixture of 1-chloro-3-methyl-2-butene (0.02 mol), 5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine] (0.015 mol),  $\text{Na}_2\text{CO}_3$  (0.015 mol) and KI (0.015 mol) in *N,N*-dimethylacetamide (150ml) was stirred at room temperature overnight. The mixture was filtered over dicalite and evaporated. The residue was taken up in  $\text{CH}_2\text{Cl}_2/\text{water}$  95/5. The precipitate was filtered off and dried. Yielding : 0.86g of 5,6-dihydro-1'-(3-methyl-2-butenyl)spiro[imidazo[2,1-b][3]benzazepine-11-[11H],4'-piperidine] monohydroiodide (12.7%); mp. 255.4°C (comp. 74).

d) A mixture of 1-(2-bromoethyl)-4-ethyl-1,4-dihydro-5H-tetrazol-5-one (0.012 mol), 5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine] (0.01 mol),  $\text{Na}_2\text{CO}_3$  (0.01 mol) and KI (10 mg) in 4-methyl-2-pentanone (200ml) was stirred and refluxed for 18 hours. The mixture was poured into water. The mixture was separated and the aqueous layer was extracted with 3-methyl-2-butanone, dried ( $\text{MgSO}_4$ ), filtered and evaporated. The residue was purified on a glass filter over silica gel (eluent :  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  95/5 to 93/7). The pure fractions were collected and evaporated. The residue was converted into the hydrochloric acid salt (1:2) in  $\text{C}_2\text{H}_5\text{OH}$ . Yielding : 2.63g of 1-[2-(5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepin-11,4'-piperidin]-1'-yl)ethyl]-4-ethyl-1,4-dihydro-5H-tetrazol-5-one dihydrochloride (56%); mp. 230°C (comp. 75).

e) A mixture of chloroacetonitrile (0.11 mol), 5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine] (0.1 mol) and *N,N*-diethylethanamine (0.12 mol) in

DMF (400ml) was stirred at room temperature for 48 hours. The mixture was poured into water and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was dried ( $\text{MgSO}_4$ ), filtered off and evaporated. The residue was purified on a glass filter over silica gel (eluent :  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  96/4). The pure fractions were collected and evaporated. The residue was crystallized from  $\text{CH}_3\text{CN}$ . Yielding : 18.5g 5,6-dihydrospiro[imidazo[2,1-b][3]-benzazepine-11-[11H],4'-piperidine]-1'-acetonitrile (63%); mp. 152.6°C (comp. 76).

#### Example B5

Bis(1,1-dimethylethyl) dicarbonate (0.095 mol) dissolved in a small amount of  $\text{CH}_2\text{Cl}_2$  was added dropwise to a stirring mixture of compound (6) (0.079 mol) in  $\text{CH}_2\text{Cl}_2$  (250ml). The mixture was stirred at room temperature for the weekend, then washed with  $\text{H}_2\text{O}$ , dried, filtered and the solvent was evaporated. Toluene was added and evaporated again. The residue was stirred in DIPE. The precipitate was filtered off and the filtrate was evaporated. This fraction was purified over silica gel on a glass filter (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0, 99/1, 98/2 and 96/4). The pure fractions were collected and the solvent was evaporated. The residue was stirred in hexane. The precipitate was filtered off and dried, yielding 15.05g of 1,1-dimethylethyl 5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-1'-carboxylate (54%) (comp. 7).

#### Example B6

a) A mixture of tetrahydro-2-furanmethanol methanesulfonate (0.01mol), compound 6 (0.01mol) and  $\text{Na}_2\text{CO}_3$  (0.02mol) in 4-methyl-2-pentanone (150ml) was stirred and refluxed overnight. The reaction mixture was filtered over dicalite. The filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  95/5). The pure fractions were collected and the solvent was evaporated. The residue was dissolved in 2-propanone and converted into the cyclohexane sulfamic acid salt (1:2). The precipitate was filtered off and dried. Yielding: 1.48g of 5,6-dihydro-1'-[(tetrahydro-2-furanyl)methyl]spiro[imidazo[2,1-b][3]benzazepine-11-[11H],4'-piperidine] cyclohexylsulfamate(1:2) monohydrate (20.7%); mp. 120.2°C (comp. 72).

b) A mixture of compound 6 (0.02 mol) and 2-thiophenecarboxaldehyde (0.053 mol) in methanol (300ml) was hydrogenated with Raney Nickel (2g) as a catalyst. After uptake of  $\text{H}_2$  (1eq), the catalyst was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent :  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  95/5). The pure fractions were collected and evaporated. The residue was converted into the cyclohexanesulfamic acid salt (1:1) in 2-propanone. The precipitate was filtered off and dried. The residue was recrystallized from 2-propanol. The precipitate was filtered off and dried. Yielding : 0.72g of 5,6-dihydro-1'-(2-thienyl-

methyl)spiro[imidazo[2,1-b][3]benzazepine-11-[11H],4'-piperidine] cyclohexyl-sulfamate(1:1) (6.6%); mp. 211.1°C (comp. 73).

#### Example B7


- 5 a) A mixture of compound (9) (0.155 mol), prepared according to the procedure described in Ex. No. B2 b, and MnO<sub>2</sub> (300g) in chloroform (1200ml) was stirred and refluxed for 90 min. The mixture was filtered over dicalite and the filtrate was evaporated. Part of this fraction (1g) was crystallized from CH<sub>3</sub>CN. The precipitate was filtered off and dried, yielding 0.5g of 1,1-dimethylethyl 3-formyl-5,6-dihydro-spiro[imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-1'-carboxylate (comp. 12).
- 10 b) A mixture of compound (12) (0.134 mol), NaCN (0.705 mol) and MnO<sub>2</sub> (233g) in methanol (2500ml) was stirred at room temperature. Acetic acid (45.5ml) was added dropwise. The mixture was stirred and refluxed for 20 hours and filtered over dicalite. The filtrate was evaporated. The residue was taken up in H<sub>2</sub>O, CH<sub>2</sub>Cl<sub>2</sub> and K<sub>2</sub>CO<sub>3</sub>. The mixture was separated into its layers. The aqueous layer was extracted with
- 15 CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layer was dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 97/3). A pure fraction was collected and the solvent was evaporated, yielding 47.7g of methyl (1,1-dimethylethyl) 5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3,1'-dicarboxylate (87%) (comp. 13).
- 20 c) A mixture of compound (13) (0.056 mol) in NaOH 1N (100ml), H<sub>2</sub>O (250ml) and THF (250ml) was stirred at room temperature for 18 hours. The organic solvent was evaporated. The aqueous concentrate was neutralized with HCl 1N (100ml) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. Part of this fraction (2g) was crystallized from CH<sub>3</sub>CN.
- 25 The precipitate was filtered off and dried, yielding 1.16g of 1'-[(1,1-dimethylethoxy)-carbonyl]-5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-carboxylic acid (comp. 14).
- d) A mixture of compound (14) (0.04 mol) and *N,N*-dimethyl-4-pyridinamine (0.04 mol) in CH<sub>2</sub>Cl<sub>2</sub> (300ml) was stirred until complete dissolution. *N,N*-diethylethanamine (0.05 mol) was added. Then *N*-(ethylcarbonimidoyl)-*N,N*-dimethyl-1,3-propanediamine monohydrochloride (0.05 mol) was added portionwise. The mixture was stirred at room temperature for 30 min. *N,N*-diethylethanamine (0.06 mol) was added and then NH<sub>4</sub>Cl (0.05 mol) was added portionwise. The mixture was stirred at
- 35 aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layer was dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was purified by column

chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  97.5/2.5). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from  $\text{CH}_3\text{CN}$ . The precipitate was filtered off and dried, yielding 9.5g of 1,1-dimethylethyl 3-(aminocarbonyl)-5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-

piperidine]-1'-carboxylate (60%) (comp. 16).

e) A mixture of compound (16) (0.023 mol) in  $\text{HCl}/2$ -propanol (25ml) and methanol (100ml) was stirred and refluxed for 90 min and then cooled. The precipitate was filtered off and dried, yielding 8g of 5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]-benzazepine-11,4'-piperidine]-3-carboxamide dihydrochloride (94%) (comp. 17). The precipitate can also be converted into the (E)-2-butenedioic acid salt.

f) Preparation of




compound 61

A mixture of compound (17) (0.0135 mol) and  $\text{NaHCO}_3$  (0.0271 mol) in THF (100ml) and ethanol (50ml) was stirred and refluxed for 10 min. Methyl 2-propenoate (0.0149 mol) was added. The mixture was stirred and refluxed for 3 hours and then cooled. The solvent was evaporated under reduced pressure. The residue was partitioned between  $\text{H}_2\text{O}$  and  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  95/5 to 90/10). The pure fractions were collected and the solvent was evaporated. The residue was purified again by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  99/1 to 97/3). The pure fractions were collected and the solvent was evaporated. The residue was refluxed in  $\text{CH}_3\text{OH}/\text{diethyl ether}$  2:8 (precipitation resulted). The precipitate was filtered off, washed with diethyl ether and dried in vacuo at  $40^\circ\text{C}$  overnight. Yielding: 2.27g of compound 61.

g) *N,N*-diethylethanamine (0.0081 mol) was added at room temperature to a suspension of compound 17 (0.004 mol) in methanol (100ml). The mixture was cooled to  $0^\circ\text{C}$ . Oxirane was bubbled through the mixture for 45 min. The mixture was allowed to warm to room temperature and then stirred at room temperature for 30 min. The solvent was evaporated under reduced pressure. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  97/3 to 95/5). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$  1:4. The precipitate was filtered off and dried. Yielding: 0.25g of 6,11-dihydro-1'-(2-hydroxyethyl)spiro[5*H*-imidazo[2,1-*b*][3]-benzazepine-11,4'-piperidine]-3-carboxamide (18%) (compound 62).



h) Preparation of




compound 63

*N,N*-diethylethanamine (0.0113 mol) was added at room temperature to a suspension of compound 17 (0.0054 mol) in methanol (100ml). After 5 min, oxirane was bubbled through the mixture at 0°C for 1 hour. The solvent was evaporated. The residue was suspended in *t*-butanol (200ml). Methyl chloroacetate (0.007 mol) and *N,N*-diethylethanamine (0.0054 mol) were added. The mixture was stirred and refluxed for 48 hours. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 98/2 to 95/5). Two fractions were collected and the solvent was evaporated. Fraction 1 was crystallized from CH<sub>3</sub>OH/CH<sub>3</sub>CN 1:3. The precipitate was filtered off and dried. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 97/3 to 95/5). The pure fractions were collected and the solvent was evaporated. Yielding: 0.46g of compound 63.

i) *N,N*-diethylethanamine (0.0271 mol) was added at room temperature to a suspension of compound 17 (0.0129 mol) and ethyl  $\alpha$ -methylenebenzeneacetate (0.0140 mol) in DMF (100ml). The mixture was stirred at room temperature over the weekend. The solvent was evaporated. The residue was extracted with CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O. The mixture was separated into its layers. The precipitate in the organic layer was filtered off. Yielding: 2.6g of ethyl 3-(aminocarbonyl)-6,11-dihydro- $\alpha$ -phenylspiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-propanoate monohydrochloride (compound 64).

j) Preparation of




compound 65

A mixture of compound 17 (0.0027 mol), 1-(3-chloropropyl)-1,3-dihydro-2*H*-benzimidazol-2-one (0.003 mol), Na<sub>2</sub>CO<sub>3</sub> (0.0027 mol) and KI (few crystals) in CH<sub>3</sub>CN (100ml) was stirred and refluxed for 48 hours. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 99/1 to 95/5). Two fractions were collected and their solvents were evaporated. The desired fraction was purified again by column chromatography over

silica gel (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  97/3 to 95/5). The pure fractions were collected and the solvent was evaporated. Yielding: 0.18g of compound 65.


k) Preparation of



compound 66

- 5 A mixture of isobutyl (2-chloroethyl)carbamate (0.008 mol), 5,6-dihydrospiro[11H-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-carboxamide dihydrochloride (0.004 mol), 4-methyl-2-pentanone (50 ml),  $\text{Na}_2\text{CO}_3$  (0.020 mol) and KI (catalytic quantity) was stirred and refluxed (oil bath:  $130^\circ\text{C}$ ) overnight. The solvent was evaporated (vacuum,  $60^\circ\text{C}$ ). Water was added.  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  90/10 was added. The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  95/5). The desired fractions were collected and the solvent was evaporated. Yield: 1.5 g of compound 66 (85.4%).

l) Preparation of



compound 67

- 15 A mixture of compound 66 (0.00227 mol) and  $\text{HCl}/2$ -propanol (3 ml) in 2-propanol (30 ml) was stirred at  $80^\circ\text{C}$  (oil bath). The solvent was evaporated (vacuum,  $40^\circ\text{C}$ ). 2-Propanol was added, then evaporated (2 x). Ethanol was added, then evaporated. The residue was stirred in boiling ethanol (50 ml), then filtered off over a P4 glass filter and the product was dried under a stream of  $\text{N}_2$ . Yield: 0.254 g (24.0%). The filtrate was stirred for 3 hours while cooling on an ice-bath. The precipitate was filtered off over a P3 glass filter and dried (vacuum,  $60^\circ\text{C}$ , 3 hours). Yield: 0.304 g (28.7%). The filtrate was evaporated. The residue was dried (over the weekend under  $\text{N}_2$  flow). Total yield: 70.4% of compound 67.

- 20 m) Methyl chloroformate (0.0036 mol) was added at room temperature to a suspension of compound (17) (0.0032 mol) in  $\text{CH}_2\text{Cl}_2$  (100ml). A mixture of *N,N*-diethylethanamine (0.0097 mol) in  $\text{CH}_2\text{Cl}_2$  (20ml) was added dropwise. The mixture was stirred at room temperature for the weekend. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  99/1 to 98/2). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from  $\text{CH}_3\text{CN}$ . The precipitate was filtered off and dried. Yield: 0.68g

of 3-(aminocarbonyl)-6,11-dihydrospiro[5*H*-imidazo[2,1-*b*][3]-benzazepine-11,4'-piperidine]-1'-carboxylate (60%) (comp. 79).

- n) A mixture of compound (17) (0.005 mol), KOAc (2g) and paraformaldehyd (0.5g) in methanol (100ml) was hydrogenated with palladium on charcoal 10% (0.5g) as a catalyst in the presence of a thiophene solution (1ml). After uptake of H<sub>2</sub> (1 equiv), the catalyst was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 97/3). The pure fractions were collected and the solvent was evaporated. This fraction was purified again by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 97/3 to 95/5). The pure fractions were collected and the solvent was evaporated. Yield: 0.43g of 6,11-dihydro-1'-methylspiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-3-carboxamide monohydrate (28%) (comp. 80).

#### Example B8

- a) A solution of compound (7) (1.63 mol) in CH<sub>2</sub>Cl<sub>2</sub> (7500ml) was cooled to 0°C under N<sub>2</sub> flow. 1-Bromo-2,5-pyrrolidinedione (1.63 mol) was added portionwise (29g each). H<sub>2</sub>O (3000ml) was added. The mixture was stirred overnight. The organic layer was separated, dried, filtered and the solvent was evaporated. This fraction was purified by HPLC over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 98/2, 90/10 and 100/0). A pure fraction was collected and the solvent was evaporated, yielding 189g of 1,1-dimethylethyl 2,3-dibromo-5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-carboxylate (27%) (comp. 48). The 3-monobromo analogue (comp. 60; Ex. No. B10a) can be prepared in a similar way.
- b) A mixture of compound (48) (0.02 mol), acetic acid, palladium(2+) salt (0.15g) and 1,3-propanediylbis[diphenylphosphine] (0.55g) in THF (150ml) was stirred in an autoclave at 150°C for 16 hours under pressure of CO gas (30 bar) and NH<sub>3</sub> gas (10 atm). The mixture was cooled, filtered and the filtrate was evaporated. This fraction was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from CH<sub>3</sub>CN. The precipitate was filtered off and dried, yielding 1,1-dimethylethyl 2,3-bis(aminocarbonyl)-5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-carboxylate (comp. 49).

#### Example B9

- Dibenzoyl peroxide (0.5g) was added to a stirring mixture of compound (7) (0.039 mol) in CH<sub>2</sub>Cl<sub>2</sub> (210ml). 1-Chloro-2,5-pyrrolidinedione (0.078 mol) in a small amount of CH<sub>2</sub>Cl<sub>2</sub> was added dropwise. The mixture was stirred at room temperature overnight. The solvent was evaporated. H<sub>2</sub>O was added and the mixture was extracted with

CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 99/1, 98/2, 96/4 and 94/6). The pure fractions were collected and the solvent was evaporated. Some starting material (7.5g; 0.02 mol) was recuperated. The reaction was carried out again. Dibenzoyl peroxide (0.5g) was added to a stirring mixture of compound (7) (0.02 mol) in CH<sub>2</sub>Cl<sub>2</sub> (210ml). 1-Chloro-2,5-pyrrolidinedione (0.078 mol) in a small amount of CH<sub>2</sub>Cl<sub>2</sub> was added dropwise. The mixture was stirred at room temperature overnight. The solvent was evaporated. H<sub>2</sub>O was added and the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 99/1 and 98.5/1.5). The pure fractions were collected and the solvent was evaporated. The residue was combined with the one obtained from the first reaction, yielding 14g of 1,1-dimethylethyl 3-chloro-5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-carboxylate (93%) (comp. 19).

#### Example B10

a) CH<sub>2</sub>Cl<sub>2</sub> (87 ml) was added to 1,1-dimethylethyl 5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-carboxylate (comp. 7) in methanol (0,0582 mol). Water (58 ml) was added. Water (58 ml) and Na<sub>2</sub>CO<sub>3</sub> (0.0582 mol) were added to the separated organic layer and the mixture was cooled to 0-5°C (solution 1).

Br<sub>2</sub> (0.0565 mol) was added to a solution of tetrabutylammoniumbromide (0.0565 mol) and CH<sub>2</sub>Cl<sub>2</sub> (29 ml). This mixture was stirred for 25 minutes at 15-25°C and added to solution 1 during 1 hour. After stirring for 1 hour at 20°C, water (58 ml) was added. The separated organic layer was evaporated. 3-Methyl-2-butanone (87 ml) and water (29 ml) were added to the oily residue and this mixture was heated to 80°C. The separated organic layer was washed with water (29ml) at 80°C. The organic layer was azeotroped till 116°C. 3-Methyl-2-butanone (40.7 ml) was distilled off and the product was crystallized during 2 hours at 50°C. The crystallized product was filtered off, washed with 3-methyl-2-butanone and dried (vacuum, 50°C). Yield : 13.03g of 1,1-dimethylethyl 3-bromo-5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-carboxylate (51.8%) (comp. 60).

b) A mixture of compound (60) (0.257 mol), DMF (515 ml), H<sub>2</sub>O (26 ml) and CuCN (1,287 mol) were heated till 132°C, stirred for 17 hours and then cooled to room temperature. The mixture was poured into 2056 ml of H<sub>2</sub>O and stirred for 2 hours. The precipitate was filtered off, washed twice with H<sub>2</sub>O (149 ml) and dried (vacuum, 100°C). The resulting precipitate was refluxed in 3-methyl-2-butanone (772 ml) for 30

min, followed by cooling the reaction mixture to 50°C and filtering. NH<sub>4</sub>OH (129 ml) was added to the filtrate at 50°C and stirred for 30 min. The 3-methyl-2-butanone layer was separated and washed with NH<sub>4</sub>OH (129 ml) as described above. This procedure was repeated for another 5 times. The 3-methyl-2-butanone layer was separated again, azeotroped for 30 min and partially evaporated. The resulting mixture was crystallized and the precipitate was filtered, washed with 3-methyl-2-butanone (7.7 ml) and dried (vacuum, 50°C). Yield : 52.4 g of 1,1-dimethylethyl 3-(aminocarbonyl)-5,6-dihydro-spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-1'-carboxylate (53%) (comp. 16).

#### 10 Example B11

a) A mixture of intermediate (18) (0.152 mol) in trifluoromethanesulfonic acid (500ml) was stirred at 158°C for 90 hours. The mixture was cooled, poured out on ice and K<sub>2</sub>CO<sub>3</sub> (800g) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated partially until 100ml while the temperature was kept below 40°C. The concentrate was purified immediately by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 95/5). The pure fractions were collected and the solvent was evaporated, yielding 18.1g of 1-methyl-spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidin]-6(5*H*)-one (42%) (comp. 22). Part of this fraction (1.5g) was dissolved in ethanol and converted into the (E)-2-butenedioic acid salt (2:3). The precipitate was filtered off and dried, yielding 1.92g of 1-methylspiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidin]-6(5*H*)-one (E)-2-butenedioic acid salt (2:3) (comp. 22a).

b) A mixture of intermediate (24) (0.041 mol) in HBr 48% solution (250ml) was stirred and refluxed for 4 hours. The mixture was poured out on ice and K<sub>2</sub>CO<sub>3</sub> and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated, yielding 10.4g of spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidin]-6(5*H*)-one (95%) (comp. 23). Part of this fraction (0.9g) was dissolved in ethanol and converted into the (E)-2-butenedioic acid salt (2:3). The precipitate was filtered off and dried, yielding 0.78g of spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidin]-6(5*H*)-one (E)-2-butenedioic acid salt (2:3) (comp. 23a).

c) A mixture of compound (23) (0.01 mol) in methanol (300ml) was stirred on an ice bath. NaBH<sub>4</sub> (0.02 mol) was added portionwise over a 15-min period. The mixture was stirred on an ice bath for 1 hour. The solvent was evaporated at a temperature below 40°C. The residue was taken up in H<sub>2</sub>O and the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 90/10. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the

solvent was evaporated, yielding 2g of 5,6-dihydrospiro[11*H*-imidazo[2,1-*b*][3]-benzazepine-11,4'-piperidine]-6-ol (75%) (comp. 26).

d) A mixture of compound (26) (0.0075 mol) in methanesulfonic acid (50ml) was stirred at room temperature for 40 min. The mixture was poured out on ice, alkalized with a NaOH 50% solution and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated, yielding 2g of spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine] (100%) (comp.27). Part of this fraction (0.3g) was dissolved in ethanol and converted into the (E)-2-butenedioic acid salt (1:1). The precipitate was filtered off and dried, yielding 0.26g spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine] (E)-2-butenedioic acid salt (1:1) (comp. 27a).

#### Example B12

A mixture of compound (24) (0.0128 mol) in H<sub>2</sub>SO<sub>4</sub> (5ml) and methanol (100ml) was stirred and refluxed for the weekend. The solvent was evaporated. The residue was taken up in H<sub>2</sub>O. The mixture was alkalized with a NaOH solution and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried, filtered and the solvent was evaporated, yielding 4.4g of 5,6-dihydro-2,3-bis(methoxymethyl)spiro[11*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine] (100%) (comp. 31).

#### Example B13


a) A mixture of 5,6-dihydrospiro[imidazo[2,1-*b*][3]benzazepine-11-[11*H*],4'-piperidine]-1'-acetonitrile (comp. 76) (0.055 mol) in NH<sub>3</sub>/CH<sub>3</sub>OH (500ml) was hydrogenated with Raney Nickel (2g) as a catalyst at room temperature. After uptake of H<sub>2</sub> (2eq), the catalyst was filtered off and the filtrate was evaporated. Yielding : 20.9g of 5,6-dihydrospiro[imidazo[2,1-*b*][3]benzazepine-11-[11*H*],4'-piperidine]-1'-ethanamine 2-propanolate(2:1). trihydrochloride. sesquihydrate; mp. 245.9°C (comp. 77).

b) A mixture of 2-chloropyrimidine (0.012mol), 5,6-dihydro-spiro[imidazo[2,1-*b*][3]-benzazepine-11-[11*H*],4'-piperidine]-1'-ethanamine (0.01mol) and Na<sub>2</sub>CO<sub>3</sub> (0.02mol) in 4-methyl-2-pentanone (200ml) was stirred and refluxed for 48 hours. The reaction mixture was filtered over dicalite. The filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 95/5). The pure fractions were collected and the solvent was evaporated. The residue was dissolved in 2-propanol and converted into the hydrochloric acid salt (1:3). The precipitate was filtered off and dried. Yielding: 0.94g of 5,6-dihydro-*N*-2-pyrimidinyl-spiro[imidazo[2,1-*b*][3]benzazepine-11-[11*H*],4'-piperidine]-1'-ethanamine trihydrochloride. monohydrate.2-propanolate(1:1) (16.7%) (comp. 78).

c) 3-Chloro-6-(3-methyl-1,2,4-thiadiazol-5-yl)pyridazine (0.01 mol) and 5,6-dihydro-spiro[imidazo[2,1-b][3]benzazepine-11-[1*H*],4'-piperidine]-1'-ethanamine (0.01 mol) were stirred at 140°C for 2h. The mixture was cooled and purified by column chromatography over silica gel (eluent : CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 95/5). The pure fractions were collected and evaporated. The residue was converted into the hydrochloric acid salt (1:4) in 2-propanol and dried. Yielding : 2.22g of 2-[[2-(5,6-dihydro-spiro[imidazo[2,1-b][3]benzazepine-11-[1*H*],4'-piperidin]-1'-yl)ethyl]amino]-4(1*H*)-pyrimidinone trihydrochloride.2-propanolate(1:1). sesquihydrate (36.6%) (comp. 68).

- 10 The following Tables list compounds of formula (I) as prepared according to one of the above examples (Ex. No.).

Table 2




Co. No.	Ex. No.	R <sub>1</sub>	R <sub>2</sub>	L	Salt / Melting point
1	B3	H	H	CH <sub>3</sub>	(1); mp. 182.8°C
3	B4a	H	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	mp. 268.5°C
4	B2a	H	H	C(=O)OCH <sub>2</sub> CH <sub>3</sub>	-
5	B2b	CH <sub>2</sub> OH	H	C(=O)OCH <sub>2</sub> CH <sub>3</sub>	-
6	B1	H	H	H	-
6a	B1	H	H	H	(2); mp. 278.5°C
7	B5	H	H	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
9	B2b	CH <sub>2</sub> OH	H	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
10	B2b	CH <sub>2</sub> OH	CH <sub>2</sub> OH	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
12	B7a	C(=O)H	H	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
13	B7b	C(=O)OCH <sub>3</sub>	H	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
14	B7c	C(=O)OH	H	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
15	B7e	C(=O)OCH <sub>3</sub>	H	H	-
15a	B7e	C(=O)OCH <sub>3</sub>	H	H	(3); -
18	B2c/B7e	CH <sub>2</sub> OH	H	H	-
18a	B2c	CH <sub>2</sub> OH	H	H	(4); -

-47-

Co. No.	Ex. No.	R <sub>1</sub>	R <sub>2</sub>	L	Salt / Melting point
19	B9	Cl	H	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
20	B7e	Cl	H	H	(3); -
24	B7e	CH <sub>2</sub> OH	CH <sub>2</sub> OH	H	-
31	B12	CH <sub>2</sub> OCH <sub>3</sub>	CH <sub>2</sub> OCH <sub>3</sub>	H	-
35	B1	CH <sub>2</sub> NHC(=O)CH <sub>3</sub>	H	H	-
39	B1	CH <sub>2</sub> NHC(=O)NH <sub>2</sub>	H	H	-
43	B1	CH <sub>2</sub> NHSO <sub>2</sub> CH <sub>3</sub>	H	H	-
48	B8a	Br	Br	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
49	B8b	C(=O)NH <sub>2</sub>	C(=O)NH <sub>2</sub>	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
51	B7e	CH <sub>2</sub> OCH <sub>3</sub>	CH <sub>2</sub> OH	H	-
52	B7e	CH <sub>2</sub> OH	CH <sub>2</sub> OCH <sub>3</sub>	H	-
53	B7e	C(=O)NH <sub>2</sub>	C(=O)NH <sub>2</sub>	H	(2); -

(1) cyclohexylsulfamate (1:2); (2) hydrochloric acid (1:2); (3) (E)-2-butenedioate (1:1); (4) (E)-2-butenedioate (2:1)

Table 3



Co No.	Ex. No.	R <sub>1</sub>	-A-B-	Salt/Melting point
8	B1	H	-CH=CF-CH=CH-	(5); -
11	B1	H	-CH=CH-CH=CCH <sub>3</sub> -	-
21	B1	H	-CH=C(OH)-CH=CH-	-
29	B1	H	-C(OH)=CH-CH=CH-	-
30	B1	H	-CH=CH-C(OH)=CH-	-
32	B1	H	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	-
32a	B1	H	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	(6); -
34	B1	H	-CH=C(OH)-C(OH)=CH-	-
46	B7e	CH <sub>2</sub> OH	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	-
50	B7e	Cl	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	-
54	B2c	H	-CH=CH-S-	-
55	B1	H	-CH=CH-N(CH <sub>3</sub> )-	-




-48-

Co No.	Ex. No.	R <sub>1</sub>	-A-B-	Salt/Melting point
57	B2c	H	-S-CH=CH-	-
58	B12	-CH <sub>2</sub> -O-CH <sub>3</sub>	-CH=CH-CH=CH-	-

(3) (E)-2-butenedioate (1:1); (5) hydrochloric acid (1:1); (6) (E)-2-butenedioate (2:3)


Table 4



Co. No.	Ex. No.	n	z	R <sub>1</sub>	L	Salt / Melting point
2	B1	2	-(CH <sub>2</sub> ) <sub>2</sub> -	H	H	(5); mp. 238.6°C
22	B11a	1	-C(=O)CH <sub>2</sub> -	H	CH <sub>3</sub>	-
22a	B11a	1	-C(=O)CH <sub>2</sub> -	H	CH <sub>3</sub>	(6); -
23	B11b	1	-C(=O)CH <sub>2</sub> -	H	H	-
23a	B11b	1	-C(=O)CH <sub>2</sub> -	H	H	(6); -
25	B1	1	-CH <sub>2</sub> -	H	H	-
25a	B1	1	-CH <sub>2</sub> -	H	H	(3); -
26	B11c	1	-CHOH-CH <sub>2</sub> -	H	H	-
27	B11d	1	-CH=CH-	H	H	-
27a	B11d	1	-CH=CH-	H	H	(3); -
28	B1	1	-(CH <sub>2</sub> ) <sub>3</sub> -	H	H	-
33	B1	1	-O-CH <sub>2</sub> -	H	H	-
33a	B1	1	-O-CH <sub>2</sub> -	H	H	(3); -
36	B5	1	-(CH <sub>2</sub> ) <sub>3</sub> -	H	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
37	B2b	1	-(CH <sub>2</sub> ) <sub>3</sub> -	CH <sub>2</sub> OH	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
38	B7e	1	-(CH <sub>2</sub> ) <sub>3</sub> -	CH <sub>2</sub> OH	H	-
42	B1	1	-(CH <sub>2</sub> ) <sub>4</sub> -	H	H	-
59	B3	2	-(CH <sub>2</sub> ) <sub>2</sub> -	H	CH <sub>3</sub>	mp. 119.2°C
60	B10a	1	-(CH <sub>2</sub> ) <sub>2</sub> -	Br	-C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-

(3) (E)-2-butenedioate (1:1); (5) hydrochloric acid (1:1); (6) (E)-2-butenedioate (2:3)


Table 5












Co. No.	Ex. No.	L	Salt / Melting point
61	B7f	$-(CH_2)_2-C(=O)OCH_3$	(7)
62	B7g	$-(CH_2)_2-OH$	-
63	B7h	$-CH_2-C(=O)OCH_3$	-
64	B7i		(5)
65	B7j		(7)
66	B7k	$-(CH_2)_2-NH-C(=O)OC(CH_3)_3$	-
67	B7l	$-(CH_2)_2-NH_2$	(8)
16	B7d/B10b	$-C(=O)OC(CH_3)_3$	-
17	B7e	H	(2); mp. 275.6°C
41	B1	H	-
79	B7m	$-C(=O)OCH_3$	-
80	B7n	$-CH_3$	(7); -

(2) hydrochloric acid (1:2); (5) hydrochloric acid (1:1); (7) monohydrate; (8) hydrochloric acid (1:3) monohydrate


## 5 Table 6



Co. No.	Ex. No.	L	Salt/Melting point
69	B4d		(9); mp. 260.4°C;
70	B4d		mp. 189.0°C
71	B4b		(1); mp. 180°C;
72	B6a		(10); mp. 120.2°C;
73	B6b		(11); mp. 211.1°C
74	B4c		(12); mp. 255.4°C
75	B4d		(2); mp. 230°C
76	B4e	-CH <sub>2</sub> -CN	mp. 152.6°C
77	B13a	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	(13); mp. 245.9°C
78	B13b		(14); mp. 216.1°C
68	B13c		(15); mp. 261.3°C

(1) cyclohexylsulfamate (1:2); (2) hydrochloric acid (1:2); (9) hydrochloric acid (1:3) hydrate (2:1) ethanolate (2:1); (10) cyclohexylsulfamate (1:2) hydrate (1:2); (11) cyclohexylsulfamate (1:1); (12) hydroiodic acid (1:1); (13) hydrochloric acid (1:3) hydrate (2:3) 2-propanolate (2:1); (14) hydrochloric acid (1:3) hydrate (1:1) 2-propanolate (1:1); (15) hydrochloric acid (1:3) hydrate (2:3) 2-propanolate (1:1)

Table 7



Co. No.	Ex. No.	z	R <sub>1</sub>	-A-B-	L	Salt
40	B5	-(CH <sub>2</sub> ) <sub>2</sub> -	H	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
44	B9	-(CH <sub>2</sub> ) <sub>2</sub> -	Cl	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
44a	B7e	-(CH <sub>2</sub> ) <sub>2</sub> -	Cl	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	(3); -
45	B2b	-(CH <sub>2</sub> ) <sub>2</sub> -	CH <sub>2</sub> OH	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	C(=O)OC(CH <sub>3</sub> ) <sub>3</sub>	-
47	B1	-CH <sub>2</sub> -	H	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	H	-
47a	B1	-CH <sub>2</sub> -	H	-CH=C(OCH <sub>3</sub> )-C(OCH <sub>3</sub> )=CH-	H	(3); -
56	B2c	-CH=CH-	H	-CH=CH-S-	H	--
56a	B2c	-CH=CH-	H	-CH=CH-S-	H	(3)

(3) (E)-2-butenedioate (1:1)

### C. Pharmacological Example

The ED<sub>50</sub> values (mg/kg) in the test "Protection of Rats from Compound 48/80 induced

5 Lethality" for the compounds of formula (I) are listed in the Table below.

Compound No.	ED <sub>50</sub> (mg/kg)
1	2.5
3	2.5
17	0.04
18a	0.08
20	0.31
53	0.31
56a	2.5
58	2.5
62	0.63
64	0.31
79	0.04
80	0.63

### D. Composition Examples

10 The following formulations exemplify typical pharmaceutical compositions suitable for systemic or topical administration to warm-blooded animals in accordance with the present invention.

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I), a prodrug, an addition salt, a N-oxide, a quaternary amine or a stereochemically isomeric form thereof.

Example D1 : Oral drops

500 g of the A.I. is dissolved in 0.5 l of 2-hydroxypropanoic acid and 1.5 l of the polyethylene glycol at 60~80°C. After cooling to 30~40°C there are added 35 l of polyethylene glycol and the mixture is stirred well. Then there is added a solution of  
5 1750 g of sodium saccharin in 2.5 l of purified water and while stirring there are added 2.5 l of cocoa flavor and polyethylene glycol q.s. to a volume of 50 l, providing an oral drop solution comprising 10 mg/ml of the A.I. The resulting solution is filled into suitable containers.

Example D2: Oral solutions

10 9 g of methyl 4-hydroxybenzoate and 1 g of propyl 4-hydroxybenzoate are dissolved in 4 l of boiling purified water. In 3 l of this solution are dissolved first 10 g of 2,3-dihydroxybutanedioic acid and thereafter 20 g of the A.I. The latter solution is combined with the remaining part of the former solution and 12 l of 1,2,3-propanetriol and 3 l of sorbitol 70% solution are added thereto. 40 g of sodium saccharin are  
15 dissolved in 0.5 l of water and 2 ml of raspberry and 2 ml of gooseberry essence are added. The latter solution is combined with the former, water is added q.s. to a volume of 20 l providing an oral solution comprising 5 mg of the A.I. per teaspoonful (5 ml). The resulting solution is filled in suitable containers.

Example D3 : Capsules

20 20 g of the A.I., 6 g sodium lauryl sulfate, 56 g starch, 56 g lactose, 0.8 g colloidal silicon dioxide, and 1.2 g magnesium stearate are vigorously stirred together. The resulting mixture is subsequently filled into 1000 suitable hardened gelatin capsules, each comprising 20 mg of the A.I..

Example D4 : Film-coated tabletsPreparation of tablet core

A mixture of 100 g of the A.I., 570 g lactose and 200 g starch is mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone (Kollidon-K 90®) in about 200 ml of water. The wet powder mixture is sieved, dried and sieved again. Then there are added 100 g microcrystalline cellulose  
30 (Avicel®) and 15 g hydrogenated vegetable oil (Stereotex ®). The whole is mixed well and compressed into tablets, giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating

35 To a solution of 10 g methyl cellulose (Methocel 60 HG®) in 75 ml of denaturated ethanol there is added a solution of 5 g of ethyl cellulose (Ethocel 22 cps ®) in 150 ml of dichloromethane. Then there are added 75 ml of dichloromethane and 2.5 ml

-53-


1,2,3-propanetriol. 10 g of polyethylene glycol is molten and dissolved in 75 ml of dichloromethane. The latter solution is added to the former and then there are added 2.5 g of magnesium octadecanoate, 5 g of polyvinylpyrrolidone and 30 ml of concentrated colour suspension (Opaspray K-1-2109®) and the whole is homogenated. The

5 tablet cores are coated with the thus obtained mixture in a coating apparatus.

-54-

Claims

1. A compound of formula



a prodrug, a *N*-oxide, an addition salt, a quaternary amine or a stereochemically isomeric form thereof wherein

$R^1$  is hydrogen,  $C_{1-6}$ alkyl, halo, formyl, carboxyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylcarbonyl,  $N(R^3R^4)C(=O)-$ ,  $N(R^3R^4)C(=O)N(R^5)-$ , ethenyl substituted with carboxyl or  $C_{1-6}$ alkyloxycarbonyl, or  $C_{1-6}$ alkyl substituted with hydroxy, carboxyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $N(R^3R^4)C(=O)-$ ,  $C_{1-6}$ alkyl $C(=O)N(R^5)-$ ,  $C_{1-6}$ alkyl $S(=O)_2N(R^5)-$  or  $N(R^3R^4)C(=O)N(R^5)-$ ;

wherein each  $R^3$  and each  $R^4$  independently are hydrogen or  $C_{1-4}$ alkyl;

$R^5$  is hydrogen or hydroxy;

$R^2$  is hydrogen,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy $C_{1-6}$ alkyl,  $N(R^3R^4)C(=O)-$ , aryl or halo;

$n$  is 1 or 2;

-A-B- represents a bivalent radical of formula

-Y-CH=CH- (a-1);

-CH=CH-Y- (a-2); or

-CH=CH-CH=CH- (a-3);

wherein each hydrogen atom in the radicals (a-1) to (a-3) may independently be replaced by  $R^6$  wherein  $R^6$  is selected from  $C_{1-6}$ alkyl, halo, hydroxy,  $C_{1-6}$ alkyloxy, ethenyl substituted with carboxyl or  $C_{1-6}$ alkyloxycarbonyl, hydroxy $C_{1-6}$ alkyl, formyl, carboxyl and hydroxycarbonyl $C_{1-6}$ alkyl; each Y independently is a bivalent radical of formula -O-, -S- or -NR<sup>7</sup>-; wherein  $R^7$  is hydrogen,  $C_{1-6}$ alkyl or  $C_{1-6}$ alkylcarbonyl;

Z is a bivalent radical of formula

-(CH<sub>2</sub>)<sub>p</sub>- (b-1),

-CH=CH- (b-2),

-CH<sub>2</sub>-CHOH- (b-3),

-CH<sub>2</sub>-O- (b-4),

-CH<sub>2</sub>-C(=O)- (b-5), or

-CH<sub>2</sub>-C(=NOH)- (b-6),

provided that the bivalent radicals (b-3), (b-4), (b-5) and (b-6) are connected to the nitrogen of the imidazole ring via their -CH<sub>2</sub>- moiety;

wherein p is 1, 2, 3 or 4;

L is hydrogen; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl;

5 C<sub>1-6</sub>alkyl substituted with one or more substituents each independently selected from hydroxy, carboxyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, aryl, aryloxy, cyano or R<sup>8</sup>HN- wherein R<sup>8</sup> is hydrogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyl; or

L represents a radical of formula

$$-\text{Alk}-\text{Y}-\text{Het}^1 \quad (\text{c-1}).$$

10                    -Alk-NH-CO-Het<sup>2</sup>                    (c-2) or


-Alk-Het<sup>3</sup> (c-3) ; wherein

Alk represents C<sub>1-4</sub>alkanediyl ;

Y represents O, S or NH;

Het<sup>1</sup>, Het<sup>2</sup> and Het<sup>3</sup> each represent furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl or imidazolyl each optionally substituted with one or two C<sub>1-4</sub>alkyl substituents; pyrrolyl or pyrazolyl optionally substituted with formyl, hydroxyC<sub>1-4</sub>alkyl, hydroxycarbonyl, C<sub>1-4</sub>alkyloxycarbonyl or with one or two C<sub>1-4</sub>alkyl substituents; thiadiazolyl or oxadiazolyl optionally substituted with amino or C<sub>1-4</sub>alkyl; pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl each optionally substituted with C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyloxy, amino, hydroxy or halo; and

Het<sup>3</sup> may also represent 4,5-dihydro-5-oxo-1H-tetrazolyl substituted with C<sub>1-4</sub>alkyl, 2-oxo-3-oxazolidinyl, 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl or a radical of formula



wherein

25 A-Z represents S-CH=CH, S-CH<sub>2</sub>-CH<sub>2</sub>, S-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>, CH=CH-CH=CH, or CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>;

aryl is phenyl or phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, hydroxy, C<sub>1-4</sub>alkyl, polyhaloC<sub>1-4</sub>alkyl, cyano, aminocarbonyl, C<sub>1-4</sub>alkyloxy or polyhaloC<sub>1-4</sub>alkyloxy;

provided that 5,6-dihydrospiro[imidazo[1,2-b][3]benzazepine-11[1*H*],4'-piperidine]  
30 and pharmaceutically acceptable addition salts thereof are not included.


2. A compound according to claim 1 wherein L is hydrogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyloxycarbonyl or C<sub>1-6</sub>alkyl substituted with hydroxy, carboxyl, C<sub>1-6</sub>alkyloxy or C<sub>1-6</sub>alkyloxycarbonyl.



3. A compound according to claim 1 wherein L is C<sub>1-6</sub>alkyl substituted with aryl and C<sub>1-6</sub>alkyloxycarbonyl.
4. A compound according to any one of the preceding claims wherein -A-B- is a bivalent radical of formula -CH=CH-CH=CH- (a-3) or -CH=CH-Y- (a-2).
5. A compound according to any one of the preceding claims wherein Z is -(CH<sub>2</sub>)<sub>p</sub>- (b-1), -CH=CH- (b-2), or -CH<sub>2</sub>-O- (b-4).
6. A compound according to claims 1, 2, 4 or 5 wherein L is hydrogen, C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl, carboxyC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, or C<sub>1-6</sub>alkyloxycarbonylC<sub>1-6</sub>alkyl.
7. A compound according to any one of the preceding claims wherein R<sup>1</sup> is hydroxyC<sub>1-6</sub>alkyl, formyl, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, N(R<sup>3</sup>R<sup>4</sup>)C(=O)-, halo or hydrogen.
8. A compound according to claim 1 wherein the compound is  
5,6-dihydrospiro[11*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-carboxamide dihydrochloride;  
1'-butyl-5,6-dihydrospiro[imidazo[2,1-b][3]benzazepine-11-[11*H*],4'-piperidine];  
6,11-dihydro-1'-methylspiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine] cyclohexylsulfamate(1:2);  
6,11-dihydrospiro[5-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-methanol] (E)-2-butenedioate (2:1);  
3-chloro-6,11-dihydrospiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine] (E)-2-butenedioate (1:1);  
6,11-dihydro-3-(methoxymethyl)spiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]. (E)-2-butenedioate (1:1);  
6,11-dihydro-1'-(2-hydroxyethyl)spiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-carboxamide;  
6,11-dihydro-1'-methylspiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-3-carboxamide monohydrate;  
ethyl 3-(aminocarbonyl)-6,11-dihydro- $\alpha$ -phenylspiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-1'-propanoate monohydrochloride;  
3-(aminocarbonyl)-6,11-dihydrospiro[5*H*-imidazo[2,1-b][3]benzazepine-11,4'-piperidine]-1'-carboxylate;

spiro[10*H*-imidazo[1,2-*a*]thieno[3,2-*d*]azepine-10,4'-piperidine];  
 6,11-dihydrospiro[5*H*-imidazo[2,1-*b*][3]benzazepine-11,4'-piperidine]-2,3-  
 dicarboxamide dihydrochloride monohydrate;  
 a prodrug, a *N*-oxide, an addition salt, a quaternary amine or a stereochemically  
 isomeric form thereof.

9. A compound of formula




a prodrug, a *N*-oxide, an addition salt, a quaternary amine or a stereochemically  
 isomeric form thereof wherein L, n, -A-B-, Z, R<sup>1</sup> and R<sup>2</sup> are defined as in claim 1  
 for use as a medicine.

10. A pharmaceutical composition comprising a pharmaceutically acceptable carrier, and  
 as active ingredient a therapeutically effective amount of a compound as described in  
 any one of claims 1 to 9.

11. A process of preparing a composition as claimed in claim 10, characterized in that, a  
 pharmaceutically acceptable carrier is intimately mixed with a therapeutically  
 effective amount of a compound as described in any one claims 1 to 9.

12. A compound of formula




a *N*-oxide, an addition salt, a quaternary amine or a stereochemically isomeric form  
 thereof wherein P is a protective group and n, -A-B-, Z, R<sup>1</sup> and R<sup>2</sup> are defined as in  
 claim 1, provided that 6,11-dihydro-1'-(phenylmethyl)-5*H*-spiro[imidazo[1,2-*b*][3]-  
 benzazepine-11,4'-piperidine] (E)-2-butenedioate(1:2) is not included.

13. A compound according to claim 12 wherein P is benzyl.


14. A process of preparing a compound as claimed in claim 1, characterized by,

- a) deprotecting an intermediate of formula (II), followed optionally by derivatizing either the piperidine moiety, or the imidazole moiety, or both the piperidine moiety and the imidazole moiety




with -A-B-, Z, L, R<sup>1</sup> and R<sup>2</sup>, and n defined as in claim 1 and P being a protective group;

- b) derivatizing an intermediate of formula (II) at the imidazole moiety, leading to the formation of an intermediate of formula (II-a), followed by deprotecting the piperidine moiety, and followed optionally by derivatizing the piperidine moiety



with -A-B-, Z, L, R<sup>1</sup> and R<sup>2</sup>, and n defined as in claim 1 and P being a protective group;


- c) by cyclizing an intermediate of formula (III) in the presence of an appropriate acid, resulting in a compound of formula (I-a)



with -A-B-, L, R<sup>1</sup> and R<sup>2</sup>, and n defined as in claim 1;


and, if desired, converting compounds of formula (I) and (I-a) into each other following art-known transformations, and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or conversely, converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid; and, if desired, preparing stereochemically isomeric forms or N-oxide forms thereof.

15. A process of preparing a compound as claimed in claim 12, characterized by,
- cyclizing a compound of formula (XI) with an appropriate acid, leading to a compound of formula (II-k), followed optionally by derivatizing the imidazole moiety, leading to a compound of formula (II-a-1)



with -A-B-,  $R^1$ ,  $R^2$ ,  $n$  and  $P$  defined as in claim 13, and  $Z_1$  being a bivalent radical of formula  $-(CH_2)_p-$ , wherein  $p$  is 1,2,3 or 4.

- by reacting a tricyclic moiety of formula (XIV) with a reagent of formula (XV) under an inert atmosphere in a reaction inert solvent in the presence of a suitable base, leading to a compound of formula (II-l), followed optionally by derivatizing the imidazole moiety leading to a compound of formula (II-a-2)



with -A-B-,  $R^1$ ,  $R^2$ ,  $n$  and  $P$  defined as in claim 13,  $W_5$  being a suitable leaving group, e.g. a halo, and  $Z_2$  being a bivalent radical of formula  $-(CH_2)_p-$ , or  $-CH_2-O-$ , wherein  $p$  is 1,2,3 or 4.

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)

Approved for use through 9/30/00. OMB 0651-0032

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

**DECLARATION FOR UTILITY OR  
DESIGN  
PATENT APPLICATION  
(37 CFR 1.63)**

- ☐ Declaration Submitted with Initial Filing **OR** ☐ Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16 (e)) required)

**Attorney Docket Number** JAB 1442 -PCT-USA

**First Named Inventor**

— COMPLETE IF KNOWN

**Application Number**

**Filing Date**

**Group Art Unit**

**Examiner Name**

**As a below named inventor, I hereby declare that:**

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

ANTIHISTAMINIC SPIRO COMPOUNDS

the specification of which

- ☐ is attached hereto  
**OR**

☒ was filed on (MM/DD/YYYY) 12/15/1999 as United States Application Number or PCT International

Application Number PCT/EP99/10176 and was amended on (MM/DD/YYYY) (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
98.204.347.3	EP	12/19/1998	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

[Page 1 of 3]

**Burden Hour Statement:** This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)

Approved for use through 9/30/00. OMB 0651-0032

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

## DECLARATION — Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

☐ Customer Number  OR

☒ Registered practitioner(s) name/registration number listed below

Place Customer Number Bar Code Label here

Name	Registration Number	Name	Registration Number
Steven P. Berman	24,772	Michael Stark	32,495
Andrea L. Colby	30,194	Ellen C. Coletti	34,140
		Mary A. Appollina	34,087

☐ Additional registered practitioner(s) named on supplemental Registered Practitioner Information sheet PTO/SB/02C attached hereto.


Direct all correspondence to: ☐ Customer Number  OR ☐ Correspondence address below

Name	Philip S. Johnson				
Address	Johnson & Johnson				
Address	One Johnson & Johnson Plaza				
City	New Brunswick	State	NJ	ZIP	08933-7003
Country	USA	Telephone	(732) 524-2359	Fax	(732) 524-2808

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))		Family Name or Surname	
Frans Eduard		Janssens	
Inventor's Signature			Date
Residence: City	Bonheiden	State	Country
			Belgium
Post Office Address	Janssen Pharmaceutica N.V., Turnhoutseweg 30		
Post Office Address			
City	Beerse	State	ZIP
			2340
Country	Belgium		

☒ Additional inventors are being named on the 1 supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto

Please type a plus sign (+) inside this box → +

PTO/SB/02A (3-97)  
Approved for use through 9/30/98 OMB 0651-0032  
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE  
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

+

## DECLARATION

ADDITIONAL INVENTOR(S)  
Supplemental Sheet  
Page 3 of 3

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle [if any])

Family Name or Surname

Joseph Elisabeth

Leenaerts

Inventor's  
Signature

*toisants*

Date

27 March  
2001

Residence: City

Rijkevorsel

State

Country

Belgium

Citizenship

Belgium

Post Office Address

Janssen Pharmaceutica N.V., Turnhoutseweg 30

BEY

Post Office Address

City

Beerse

State

ZIP

2340

Country

Belgium

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle [if any])

Family Name or Surname

Inventor's  
Signature

Date

Residence: City

State

Country

Citizenship

Post Office Address

Post Office Address

City

State

ZIP

Country

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle [if any])

Family Name or Surname

Inventor's  
Signature

Date

Residence: City

State

Country

Citizenship

Post Office Address

Post Office Address

City

State

ZIP

Country

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

+

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

## In Re Application of:

Frans Eduard JANSSEN and  
Joseph Elizabeth LEENAERTS

Group Art Unit: Not yet known

International Application No. : PCT/EP99/10176

Examiner: Not yet assigned

International Filing Date: 15 December 1999

For: ANTIHISTAMINIC SPIRO  
COMPOUNDS

Assistant Commissioner for Patents  
Washington DC 20231

Sir:

## ASSOCIATE POWER OF ATTORNEY

The undersigned, of the firm WOODCOCK WASHBURN KURTZ  
MACKIEWICZ & NORRIS LLP, One Liberty Place - 46th Floor, Philadelphia, Pennsylvania  
19103, Attorney and/or Agents for Applicant(s), hereby appoints the following:


Richard E. Kurtz	Registration No. 19,263	Lynn A. Malinoski	Registration No. 38,788
Dale M. Heist	Registration No. 28,425	Steven B. Samuels	Registration No. 37,711
John W. Caldwell	Registration No. 28,937	Anthony J. Rossi	Registration No. 24,053
Gary H. Levin	Registration No. 28,734	Terence P. Strobaugh	Registration No. 25,460
Steven J. Rocci	Registration No. 30,489	Michael J. Swope	Registration No. 38,041
Dianne B. Elderkin	Registration No. 28,598	Michael J. Bonella	Registration No. 41,628
John P. Donohue, Jr.	Registration No. 29,916	Harold H. Fullmer	Registration No. 42,560
Henrik D. Parker	Registration No. 31,863	John E. McGlynn	Registration No. 42,863
Suzanne E. Miller	Registration No. 32,279	Jonathan M. Waldman	Registration No. 40,861
Lynn B. Morreale	Registration No. 32,842	Paul K. Legaard	Registration No. 38,534
Mark DeLuca	Registration No. 33,229	Chad Ziegler	Registration No. 44,273
Joseph Lucci	Registration No. 33,307	Gwilym J.O. Attwell	Registration No. 45,449
Michael P. Dunnam	Registration No. 32,611	Maureen S. Gibbons	Registration No. 44,121
Michael D. Stein	Registration No. 34,734	Steven H. Meyer	Registration No. 37,189
Albert J. Marcellino	Registration No. 34,664	Paul B. Milcetic	Registration No. 46,261
David R. Bailey	Registration No. 35,057	Joseph R. Condo	Registration No. 42,431
Doreen Yatko Trujillo	Registration No. 35,719	Michael K. Jones	Registration No. 41,100
Barbara L. Mullin	Registration No. 38,250	Frank T. Carroll	Registration No. 42,392
Michael P. Straher	Registration No. 38,325	Mark J. Rosen	Registration No. 39,822
Kevin M. Flannery	Registration No. 35,871	Mitchell R. Brustein	Registration No. 38,394
David A. Cherry	Registration No. 35,099	Eric H. Vance	Registration No. 47,151
Robert B. Washburn	Registration No. 16,574	Peter M. Ullman	Registration No. 43,963
Norman L. Norris	Registration No. 24,196	Thomas E. Watson	Registration No. 43,243



Richard B. LeBlanc	Registration No. 39,495
Joseph D. Rossi	Registration No. 47,038
Rosaleen P. Morris	Registration No. 47,321
George J. Awad	Registration No. 46,528
Steven D. Maslowski	Registration No. 46,905
S. Maurice Valla	Registration No. 43,966
Vincent J. Roccia	Registration No. 43,887
Robin S. Quartin	Registration No. 45,028
Patrick J. Farley	Registration No. 42,524
Gregory L. Hillyer	Registration No. 44,154
Susan C. Murphy	Registration No. 46,221
David L. Marcus	Registration No. 46,897
Christine A. Goddard	Registration No. 46,731
Ellen M. Klann	Registration No. 44,836
Andrew J. Hagerty	Registration No. 44,141
Emma R. Dailey	Registration No. P48,491
Stephen C. Timmins	Registration No. P48,481
Erich M. Falke	Registration No. P49,049
Jane Smith Ingelese	Registration No. P48,444

his/her associates with full power to prosecute the above-identified application and to transact all business in the Patent Office connected therewith and requests that correspondence continue to be directed to the firm of WOODCOCK WASHBURN KURTZ MACKIEWICZ & NORRIS LLP at the above address.

Date: 6/19/01



David N. Farsiou

Registration No. 44,104

Woodcock Washburn Kurtz  
Mackiewicz & Norris LLP  
One Liberty Place - 46th Floor  
Philadelphia PA 19103  
Telephone: (215) 568-3100  
Facsimile: (215) 568-3439

DOCKET NO. JANS-0004

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re Application of:

Frans Eduard JANSSEN and  
Joseph Elizabeth LEENAERTS

Group Art Unit: Not yet known

International Application No. : PCT/EP99/10176

Examiner: Not yet assigned

International Filing Date: 15 December 1999

For: ANTIHISTAMINIC SPIRO  
COMPOUNDS

Assistant Commissioner for Patents  
Washington DC 20231

Sir:

ASSOCIATE POWER OF ATTORNEY


The undersigned, Mary A. Appollina, Registration No. 34,087, of **Johnson & Johnson**, One Johnson Plaza, New Brunswick, N.J. 08933-7002, hereby appoints the following of the firm **WOODCOCK WASHBURN KURTZ MACKIEWICZ & NORRIS LLP**, One Liberty Place - 46th Floor, Philadelphia, Pennsylvania 19103, Attorney and/or Agents for Applicant(s):

Richard E. Kurtz	Registration No. 19,263	David A. Cherry	Registration No. 35,099
Dale M. Heist	Registration No. 28,425	Robert B. Washburn	Registration No. 16,574
John W. Caldwell	Registration No. 28,937	Norman L. Norris	Registration No. 24,196
Gary H. Levin	Registration No. 28,734	Lynn A. Malinoski	Registration No. 38,788
Steven J. Rocci	Registration No. 30,489	Steven B. Samuels	Registration No. 37,711
Dianne B. Elderkin	Registration No. 28,598	Anthony J. Rossi	Registration No. 24,053
John P. Donohue, Jr.	Registration No. 29,916	Terence P. Strobaugh	Registration No. 25,460
Henrik D. Parker	Registration No. 31,863	Michael J. Swope	Registration No. 38,041
Suzanne E. Miller	Registration No. 32,279	Michael J. Bonella	Registration No. 41,628
Lynn B. Morreale	Registration No. 32,842	Harold H. Fullmer	Registration No. 42,560
Mark DeLuca	Registration No. 33,229	John E. McGlynn	Registration No. 42,863
Joseph Lucci	Registration No. 33,307	Jonathan M. Waldman	Registration No. 40,861
Michael P. Dunnam	Registration No. 32,611	Paul K. Legaard	Registration No. 38,534
Michael D. Stein	Registration No. 34,734	Chad Ziegler	Registration No. 44,273
Albert J. Marcellino	Registration No. 34,664	Gwilym J.O. Attwell	Registration No. 45,449
David R. Bailey	Registration No. 35,057	David N. Farsiou	Registration No. 44,104
Doreen Yatko Trujillo	Registration No. 35,719	Maureen S. Gibbons	Registration No. 44,121
Barbara L. Mullin	Registration No. 38,250	Steven H. Meyer	Registration No. 37,189
Michael P. Straher	Registration No. 38,325	Paul B. Milcetic	Registration No. 46,261
Kevin M. Flannery	Registration No. 35,871	Joseph R. Condo	Registration No. 42,431

Michael K. Jones	Registration No. 41,100
Frank T. Carroll	Registration No. 42,392
Mark J. Rosen	Registration No. 39,822
Mitchell R. Brustein	Registration No. 38,394
Eric H. Vance	Registration No. 47,151
Peter M. Ullman	Registration No. 43,963
Thomas E. Watson	Registration No. 43,243
Richard B. LeBlanc	Registration No. 39,495
Joseph D. Rossi	Registration No. 47,038
Rosaleen P. Morris	Registration No. 47,321
George J. Awad	Registration No. 46,528
Steven D. Maslowski	Registration No. 46,905
S. Maurice Valla	Registration No. 43,966
Vincent J. Roccia	Registration No. 43,887
Robin S. Quartin	Registration No. 45,028
Patrick J. Farley	Registration No. 42,524
Gregory L. Hillyer	Registration No. 44,154
Susan C. Murphy	Registration No. 46,221
David L. Marcus	Registration No. 46,897
Christine A. Goddard	Registration No. 46,731
Ellen M. Klann	Registration No. 44,836
Andrew J. Hagerty	Registration No. 44,141
Emma R. Dailey	Registration No. P48,491
Stephen C. Timmins	Registration No. P48,481
Erich M. Falke	Registration No. P49,049
Jane Smith Ingelese	Registration No. P48,444

his/her associates with full power to prosecute the above-identified application and to transact all business in the Patent Office connected therewith and requests that correspondence continue to be directed to the firm of **WOODCOCK WASHBURN KURTZ MACKIEWICZ & NORRIS LLP** at One Liberty Place - 46th Floor, Philadelphia PA 19103; Telephone: (215) 568-3100; Facsimile: (215) 568-3439.

Date: *July 11, 2001*

  
Mary A. Appollina  
Registration No.: 34,087  
Johnson & Johnson